

Physicists find evidence of exotic charge transport in quantum material

November 23 2023



Credit: Pixabay/CC0 Public Domain

True to form, a "strange metal" quantum material proved strangely quiet in recent quantum noise experiments at Rice University. [Published this week](#) in *Science*, the measurements of quantum charge fluctuations known as "shot noise" provide the first direct evidence that electricity seems to flow through strange metals in an unusual liquidlike form that

cannot be readily explained in terms of quantized packets of charge known as quasiparticles.

"The noise is greatly suppressed compared to ordinary wires," said Rice's Doug Natelson, the study's corresponding author. "Maybe this is evidence that quasiparticles are not well-defined things or that they're just not there and charge moves in more complicated ways. We have to find the right vocabulary to talk about how charge can move collectively."

The experiments were performed on nanoscale wires of a well-studied quantum critical material with a precise 1-2-2 ratio of ytterbium, rhodium and silicon (YbRh_2Si_2). The material contains a high degree of quantum entanglement that produces temperature-dependent behavior.

If cooled below a [critical temperature](#), for example, the material instantly transitions from non-magnetic to magnetic. At temperatures slightly above the critical threshold, YbRh_2Si_2 is a "heavy fermion" metal, with charge-carrying quasiparticles that are hundreds of times more massive than bare electrons.

In metals, each quasiparticle, or discrete unit, of charge is the product of incalculable tiny interactions between countless electrons. First put forward 67 years ago, the quasiparticle is a concept physicists use to represent the combined effect of those interactions as a single quantum object for the purposes of quantum mechanical calculations.

Some prior theoretical studies have suggested that strange metal charge carriers might not be quasiparticles, and shot noise experiments allowed Natelson, study lead author Liyang Chen, a former student in Natelson's lab, and more than a dozen co-authors from Rice and the Technical University of Vienna (TU-Wien) to gather the first direct empirical evidence to test the idea.

"The shot noise measurement is basically a way of seeing how granular the charge is as it goes through something," said Natelson, professor of physics and astronomy, electrical and computer engineering and materials science and nanoengineering.

"The idea is that if I'm driving a current, it consists of a bunch of discrete charge carriers. Those arrive at an average rate, but sometimes they happen to be closer together in time, and sometimes they're farther apart."

Applying the technique in crystals made from the 1-2-2 ratio of ytterbium, rhodium and silicon presented significant technical challenges. For example, the crystalline films, which were grown in the laboratory of lead TU-Wien co-author Silke Paschen, had to be nearly perfect. And Chen had to find a way to maintain that level of perfection while fashioning wires from the crystal that were about 5,000 times narrower than a human hair.

Rice co-author Qimiao Si, the lead theorist on the study and the Harry C. and Olga K. Wiess Professor of Physics and Astronomy, said he, Natelson and Paschen first discussed the idea for the experiments while Paschen was a visiting scholar at Rice in 2016. Si said the results are consistent with a theory of quantum criticality he published in 2001 and has continued to explore in a nearly two-decade collaboration with Paschen.

"The low shot noise brought about fresh new insights into how the charge-current carriers entwine with the other agents of the quantum criticality that underlies the strange metallicity," said Si, whose group performed calculations that ruled out the [quasiparticle](#) picture. "In this theory of quantum criticality, the electrons are pushed to the verge of localization, and the quasiparticles are lost everywhere on the Fermi surface."

Natelson said the larger question is whether similar behavior might arise in any or all of the dozens of other compounds that exhibit strange metal behavior.

"Sometimes you kind of feel like nature is telling you something," Natelson said. "This 'strange metallicity' shows up in many different physical systems despite the fact that the microscopic, underlying physics is very different. In copper-oxide superconductors, for example, the microscopic physics is very, very different than in the heavy-fermion system we're looking at. They all seem to have this linear-in-temperature resistivity that's characteristic of [strange metals](#), and you have to wonder is there something generic going on that is independent of whatever the microscopic building blocks are inside them."

More information: Liyang Chen et al, Shot noise in a strange metal, *Science* (2023). [DOI: 10.1126/science.abq6100](https://doi.org/10.1126/science.abq6100).
www.science.org/doi/10.1126/science.abq6100

Provided by Rice University

Citation: Physicists find evidence of exotic charge transport in quantum material (2023, November 23) retrieved 2 May 2024 from <https://phys.org/news/2023-11-physicists-evidence-exotic-quantum-material.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.