


Exploring the limitations of asteroid crater lakes as climate archives

December 5 2023



Drill core from borehole taken in 1981 from the Nördlinger Ries. There are distinctive layers of light-colored dolomite between fine-layered dark sediments of the former crater lake. Credit: Gernot Arp, Göttingen University


In southern Germany just north of the Danube, there lies a large circular depression between the hilly surroundings: the Nördlinger Ries. Almost 15 million years ago, an asteroid struck this spot. Today, the impact crater is one of the most useful analogs for asteroid craters on early Mars.

Studying the deposits of the former lake that formed in the crater is particularly informative. These deposits have been of great interest ever since NASA began exploring Martian craters for signs of water and life on Mars. However, the chemical development of the former [crater lake](#) and its habitable areas is only partially understood.

An international research team led by the University of Göttingen has now uncovered clues about the past: they analyzed dolomite rocks in a drill core and found an extremely high proportion of the carbon isotope C-13. Further investigations traced this back to a phase of strong methane formation by microorganisms known as archaea in water with a low sulfate content.

In contrast, the sediments of the previous, first phase of the crater lake showed clear traces of high sulfate content and bacterial sulfate decomposition. This change reveals that the groundwater pathways to the lake changed as the [crater floor](#) cooled. The results have been [published](#) in the journal *Geochimica et Cosmochimica Acta*.

A 250-meter-long drill core taken in 1981 provided information about the [chemical processes](#) during the time periods that sediment was being deposited in the crater lake. Combining sedimentological, biogeochemical and isotope geochemical research methods enabled the researchers to identify a distinctive section, which they investigated in more detail using biomarker analyses.



Scanning electron microscope image using material taken in the drill core from the borehole taken in 1981 from the Nördlinger Ries reveals dolomite crystals with rhombohedral shape. Credit: Lingqi Zeng, Göttingen University

They detected organic biomarkers originating from sulfate-reducing bacteria and "normal" dolomite in older rocks from the crater lake. In the younger rocks, they found dolomite enriched with C-13 and a chemical called archaeol which indicates that archaea were present at that time.

The properties of the rocks reflect the conditions in the crater lake during their formation: the decrease in sulfate is due to degradation by bacteria and the C-13 enrichment is due to the formation of methane by

archaea.

"This chemical development can only be explained by the change in the groundwater supply during the gradual cooling of the crater floor. This led to a change from deep, hydrothermal groundwater (with sulfate) to cooler water without sulfate that must have flowed through limestone rocks near to the surface," explains study leader Professor Gernot Arp from the Department of Geobiology at the University of Göttingen.

The findings not only provide important information on the development of the crater lake being investigated, but also, as Arp notes, "Our findings show that the conditions in asteroid crater lakes are strongly controlled by internal processes such as crater floor cooling and water supply. In contrast, [climatic changes](#) are of secondary importance, unlike in many other lakes. This must be taken into account when deposits in terrestrial and extraterrestrial craters are used as climate archives to deduce past climate conditions from the sediments."

More information: Lingqi Zeng et al, Extremely ^{13}C -enriched dolomite records interval of strong methanogenesis following a sulfate decline in the Miocene Ries impact crater lake, *Geochimica et Cosmochimica Acta* (2023). [DOI: 10.1016/j.gca.2023.10.013](https://doi.org/10.1016/j.gca.2023.10.013)

Provided by University of Göttingen

Citation: Exploring the limitations of asteroid crater lakes as climate archives (2023, December 5) retrieved 14 May 2024 from <https://phys.org/news/2023-12-exploring-limitations-asteroid-crater-lakes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.