


From infamy to ingenuity—bacterial hijack mechanisms as advanced genetic tools

December 5 2023



Crystal structures of SAP05–SPL5^{ZnF} and SAP05–Rpn10^{vWA} complexes revealing the bimodular architecture of SAP05. (A) Cartoon model illustrating the fold of the SAP05 effector. α -helices, β -sheets, and loops are indicated in cyan, red, and blue, respectively. (B) Amino acid sequence of SAP05 highlighting the locations of secondary structures (shown in A, color matched) and SPL5^{ZnF} (purple) and Rpn10^{vWA} interacting residues (orange) (shown in C–F, color-matched). SPL5^{ZnF}, zinc-finger domain of SPL5 transcription factors. Rpn10^{vWA}, von Willebrand factor type A domain of Rpn10 ubiquitin receptor.

(C) Crystal structure of the SAP05–SPL5^{ZnF} complex (PDB 8PFC). The Zn²⁺ ions bound to the two ZnF domains are shown in gray. (D) Crystal structure of the SAP05–Rpn10^{vWA} complex (PDB 8PFD). In (C) and (D), the dashed lines indicate the interactions between the residues from both components. (E) Interfaces of SAP05 showing the loop surface (purple) and sheet surface (orange) residues that bind to SPL5^{ZnF} and Rpn10^{vWA}, respectively. (F) Gel filtration chromatograms of SAP05, SPL5^{ZnF}, and Rpn10^{vWA} and binary and ternary complexes of these proteins. The Coomassie-stained protein SDS-PAGE gel shows the presence of the three proteins in the SPL5^{ZnF}–SAP05–Rpn10^{vWA} ternary complex—see [SI Appendix, Fig. S1](#), for the other complexes. Elution volumes are indicated at the bottom of each peak with the same colors. (G) Hypothetical ternary structure of SPL5^{ZnF}–SAP05–Rpn10^{vWA} obtained by superimposing the crystal structures of SAP05–SPL5^{ZnF} and SAP05–Rpn10^{vWA} complexes. Credit: *Proceedings of the National Academy of Sciences* (2023). DOI: 10.1073/pnas.2310664120

Researchers have uncovered the intricate molecular mechanism used by parasitic phytoplasma bacteria, known for inducing "zombie-like" effects in plants. This detailed revelation opens new horizons for groundbreaking applications in biotechnology and even in biomedicine.

The team led by Professor Saskia Hogenhout at the John Innes Centre, in partnership with The Sainsbury Laboratory, has employed X-ray crystallography to unveil the structure and functional mechanism of SAP05. This molecule plays a crucial role in bridging two distinct components inside plant cells.

The discovery sheds new light on a peculiar phenomenon in nature—mostly seen in "witches' brooms" in which plant stems and leaves proliferate due to Phytoplasma bacteria.

This insect-transmitted bacteria triggers diseases like Aster Yellows,

significantly diminishing yields in leaf crops including [oilseed rape](#), lettuce, carrots, grapevines, onions, and a variety of ornamental and vegetable crops worldwide.

[Previous research](#) by the Hogenhout group revealed how the bacterial [protein](#) SAP05 is able to manipulate plants by hijacking molecular machinery called the proteasome. The proteasome breaks down and recycles proteins that are no longer required inside plant cells.

SAP05 hijacks this process, causing proteins that regulate growth and development to be dispensed into a molecular recycling center known as the 26S proteasome.

This latest research focuses on how this happens at the structural level. SAP05 effectively disrupts the molecular recycling pathway, serving as a scaffold that connects its two cellular targets: a transcription factor and the proteasome.

Fascinatingly, SAP05 binds in a manner that enables it to 'lift the dustbin lid,' selectively disposing of developmental proteins, while strategically preserving functions vital for the survival of its plant host.

Professor Hogenhout, group leader at the John Innes Center and lead of the research team behind these findings, explains, "We now know the structure of this complex and how the protein binds to two cellular components to create a [short circuit](#). While SAP05 allows itself to get involved in the plant, it does not disrupt other important processes. It is so amazing to see evolution crystallized in this way."

Usually, in plants, in fact, across all [multicellular organisms](#), this recycling of proteins in the proteasome is dependent on a molecule called ubiquitin.

By short-circuiting this process, SAP05 provides a new way of carrying out the essential task of protein degradation, which serves its own parasitic purpose and is completely independent of ubiquitin.

The discovery presents some intriguing possibilities. The researchers were struck by the sophisticated ingenuity of SAP05, a master manipulator, and its promising applications in biotechnology.

First author of the paper Dr. Qun Liu says, "It was so exciting to see that this molecule SAP05 had two sides, one side binding to the transcription factor and the other binding to the 26S [proteasome](#). It's very smart."

By understanding how this bacterial mechanism interacts with cells at a structural level, the researchers can now use this knowledge to engineer SAP05-like molecules which could be repurposed to remove unwanted proteins such as pathogen effectors or viruses, with an impact on therapeutics, research, and in agriculture.

The study is [published](#) in the journal *Proceedings of the National Academy of Sciences*.

More information: Qun Liu et al, Bimodular architecture of bacterial effector SAP05 that drives ubiquitin-independent targeted protein degradation, *Proceedings of the National Academy of Sciences* (2023).
[DOI: 10.1073/pnas.2310664120](https://doi.org/10.1073/pnas.2310664120)

Provided by John Innes Centre

Citation: From infamy to ingenuity—bacterial hijack mechanisms as advanced genetic tools (2023, December 5) retrieved 29 April 2024 from <https://phys.org/news/2023-12-infamy-ingenuitybacterial-hijack-mechanisms-advanced.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.