


# Molecular fossils study sheds light on feeding strategy shift in ancient life

December 7 2023, by Andy Fell



Overview of the Neoproterozoic sterane record. The geologic timescale is provided on the left, with dates in hundreds of millions of years. Arrows indicate the approximate time when various biomarkers become detectable above background thresholds. The site where sterols are normally methylated by the gene *24-C sterol methyltransferase* (carbon-24) is noted on ergostane with an arrow. Credit: *Nature Communications* (2023). DOI: 10.1038/s41467-023-43545-z

Paleontologists are getting a glimpse at life over a billion years in the past based on chemical traces in ancient rocks and the genetics of living animals. Research [published](#) in *Nature Communications* combines geology and genetics, showing how changes in the early Earth prompted a shift in how animals eat.

David Gold, associate professor in the Department of Earth and Planetary Sciences at the University of California, Davis, works in the new field of molecular paleontology, using geology and biology tools to study life's evolution. With new technology, it's possible to recover chemical traces of life from [ancient rocks](#), where [animal fossils](#) are scarce.

Lipids, in particular, can survive in rocks for hundreds of millions of years. Traces of sterol lipids, which come from cell membranes, have been found in rocks up to 1.6 billion years old. Currently, most [animals](#) use cholesterol—sterols with 27 carbon atoms ( $C_{27}$ )—in their cell membranes. In contrast, fungi typically use  $C_{28}$  sterols, while plants and [green algae](#) produce  $C_{29}$  sterols. The  $C_{28}$  and  $C_{29}$  sterols are also known as phytosterols.

$C_{27}$  sterols have been found in rocks 850 million years old, while  $C_{28}$  and  $C_{29}$  traces appear about 200 million years later. This is thought to reflect the increasing diversity of life at this time and the evolution of the first fungi and green algae.

Without actual fossils, it's hard to say much about the animals or plants these sterols came from. However, a [genetic analysis](#) by Gold and colleagues is shedding some light.

## Don't make it, eat it

Most animals are not able to make phytosterols themselves, but they can

obtain them by eating plants or fungi. Recently, it was discovered that annelids (segmented worms, a group that includes the common earthworm) have a gene called *smt*, which is required to make longer-chain sterols. By looking at *smt* genes from different animals, Gold and colleagues created a [family tree](#) for *smt* first within the annelids and then across animal life in general.

They found that the gene originated very far back in the evolution of the first animals and then went through rapid changes around the same time that phytosterols appeared in the rock record. Subsequently, most lineages of animals lost the *smt* gene.

"Our interpretation is that these phytosterol molecular fossils record the rise of algae in ancient oceans and that animals abandoned phytosterol production when they could easily obtain it from this increasingly abundant food source," Gold said. "If we're right, then the history of the *smt* gene chronicles a change in animal feeding strategies early in their evolution."

**More information:** T. Brunoir et al, Common origin of sterol biosynthesis points to a feeding strategy shift in Neoproterozoic animals, *Nature Communications* (2023). [DOI: 10.1038/s41467-023-43545-z](https://doi.org/10.1038/s41467-023-43545-z)

Provided by UC Davis

Citation: Molecular fossils study sheds light on feeding strategy shift in ancient life (2023, December 7) retrieved 12 May 2024 from <https://phys.org/news/2023-12-molecular-fossils-strategy-shift-ancient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---