

X-ray method enables micron-resolution imaging of living organisms over long time periods

December 7 2023



The new X-ray imaging technique uses a much lower X-ray dose thanks to two Bragg magnifier crystals (center) and a single-photon-counting detector (on the left). The sample is shown on the right. Credit: Rebecca Spiecker, Karlsruhe Institute of Technology



Researchers have developed an X-ray imaging technique that can produce detailed images of living organisms with a much lower X-ray dose than previously possible. The advance enables small organisms or other sensitive samples to be studied at high resolution over much longer periods, which could reveal new insights into a variety of dynamic processes.

The approach is based on phase contrast imaging, which relies not only on the <u>absorption</u> of X-rays in a sample, but also on the wave properties of X-rays. More precisely, it creates images from phase changes that occur as X-rays go through a specimen.

"Previously, micrometer <u>resolution</u> X-ray phase contrast imaging of living organisms was only possible for a few seconds up to minutes because severe radiation damage would occur," explained researcher team member Rebecca Spiecker from Karlsruhe Institute of Technology in Germany. "We reduced the necessary X-ray dose by overcoming the current limitations of high-resolution imaging for dose-sensitive applications."

In the journal *Optica*, the researchers describe how they developed a new X-ray imaging system that uses dedicated highly-efficient X-ray optics and single-photon-counting detectors to boost the dose efficiency for full-field imaging at micrometer resolution. They demonstrated the benefit of the new technique by imaging tiny parasitoid wasps emerging from their host eggs for more than 30 minutes.

"We show that our method exhibits superior imaging performance compared to a conventional high-resolution detector," said Spiecker. "This could be useful, for example, for capturing details of the development and behavior of small model organisms, such as Xenopus



frog embryos, over a longer time scale than is currently possible."

Better images with less radiation

X-ray imaging can reveal hidden structures and processes in living organisms. However, it also exposes organisms to radiation that is harmful at high doses, limiting how long observations can last before damage occurs. This is aggravated by the fact that the detection efficiency of commonly used high-resolution detectors decreases with increasing resolution, which means that even higher X-ray doses are necessary to obtain a high-resolution image.

To overcome this challenge, the researchers developed a phase contrast imaging approach that directly magnifies the X-ray image rather than converting the X-ray image into a visible light image and then magnifying it, which is the typical method. This allowed them to use highly efficient large-area detectors while maintaining micrometer spatial resolution.





A new X-ray imaging technique can produce detailed images of living organisms with a much lower X-ray dose than previously possible. The researchers used the new technique to image tiny parasitoid wasps emerging from their host eggs for more than 30 minutes. Credit: Rebecca Spiecker, Karlsruhe Institute of Technology

In the new imaging system, the researchers used a single-photoncounting imaging detector with a pixel size of 55 microns. The X-ray image is magnified behind the sample using crystal optics, known as a Bragg magnifier. The latter consists of two perfect silicon crystals to perform magnification.

"To achieve the highest possible dose efficiency for full-field X-ray imaging at micrometer resolution, we combine X-ray phase contrast, a Bragg magnifier and a single-photon-counting detector, all optimized for an optimal X-ray energy of 30 keV," said Spiecker. "The concept of



Bragg magnifiers dates back to the late 1970s, and although their potential for increasing dose efficiency has been noted, it has not been explored until now."

After showing that their new system could attain a dose efficiency of more than 90% while providing a resolution of up to 1.3 microns, the researchers compared its performance to a conventional high-resolution detector system using the same sample, X-ray fluence and 30 keV X-ray energy.

"At this energy, we showed that the detective quantum efficiency of our system exceeds the conventional system by over two orders of magnitude for the relevant high-resolution components of the image," said Spiecker. "This results in better images and allows a drastic reduction in the X-ray dose in the sample."

Imaging tiny insects

The researchers then used the system to perform a pilot behavioral study on living parasitoid wasps, which are widely used for biological pest control. Thanks to the minimal radiation exposure, they were able to capture images of the tiny wasps inside their host eggs for 30 minutes before the wasps finally emerged.

The researchers say that the method might also be useful for biomedical applications, such as gentle tomographic examination of biopsy samples. However, using a Bragg magnifier requires a monochromatic, coherent and collimated beam, which is available at X-ray synchrotron facilities. They are also continuing to improve the system to achieve a larger field of view and increased long-term mechanical stability for even longer measurement times.

More information: Rebecca Spiecker et al, Dose-efficient in vivo X-



ray phase contrast imaging at micrometer resolution, *Optica* (2023). DOI: 10.1364/OPTICA.500978

Provided by Optica

Citation: X-ray method enables micron-resolution imaging of living organisms over long time periods (2023, December 7) retrieved 9 May 2024 from <u>https://phys.org/news/2023-12-x-ray-method-enables-micron-resolution-imaging.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.