


# 'Giant' predator worms more than half a billion years old discovered in North Greenland

January 3 2024

---



Digital 3D model of *T. koprii* gen. et sp. nov. Reconstruction showing internal and external anatomy (red, musculature; blue, ventral ganglion; black, jaw apparatus; green, gut). (A) Lateral view. (B) Dorsal view. (C) Ventral view. (D)

Ventral view excluding musculature. Credit: *Science Advances* (2024). DOI: 10.1126/sciadv.adi6678

Fossils of a new group of animal predators have been located in the Early Cambrian Sirius Passet fossil locality in North Greenland. These large worms may be some of the earliest carnivorous animals to have colonized the water column more than 518 million years ago, revealing a past dynasty of predators that scientists didn't know existed.

The new fossil animals have been named Timorebestia, meaning 'terror beasts' in Latin. Adorned with fins down the sides of their body, a distinct head with long antennae, massive jaw structures inside their mouth, and growing to more than 30cm in length, these were some of the largest swimming animals in the Early Cambrian times.

"We have previously known that primitive arthropods were the dominant predators during the Cambrian, such as the bizarre-looking anomalocaridids," said Dr. Jakob Vinther from the University of Bristol's Schools of Earth Sciences and Biological Sciences, a senior author on the study. "However, Timorebestia is a distant, but close, relative of living arrow worms, or chaetognaths. These are much smaller ocean predators today that feed on tiny zooplankton."

"Our research shows that these ancient ocean ecosystems were fairly complex, with a food chain that allowed for several tiers of predators."

"Timorebestia were giants of their day and would have been close to the top of the [food chain](#). That makes it equivalent in importance to some of the top carnivores in modern oceans, such as sharks and seals back in the Cambrian period."

Inside the fossilized digestive system of *Timorebestia*, the researchers found remains of a common, swimming arthropod called *Isoxys*. "We can see these arthropods were a [food source](#) for many other animals," said Morten Lunde Nielsen, a former Ph.D. student at Bristol and part of the current study.

"They are very common at Sirius Passet and had long protective spines, pointing both forwards and backwards. However, they clearly didn't completely succeed in avoiding that fate, because *Timorebestia* munched on them in great quantities."

Arrow worms are one of the oldest animal fossils from the Cambrian. While arthropods appear in the [fossil record](#) about 521 to 529 million years ago, arrow worms can be traced back at least 538 million years back in time.

Dr. Vinther explained, "Both arrow worms, and the more primitive *Timorebestia*, were swimming predators. We can therefore surmise that in all likelihood they were the predators that dominated the oceans before arthropods took off. Perhaps they had a dynasty of about 10–15 million years before they got superseded by other, and more successful, groups."

Luke Parry from Oxford University, who was part of the study, added, "*Timorebestia* is a really significant find for understanding where these jawed predators came from. Today, arrow worms have menacing bristles on the outside of their heads for catching prey, whereas *Timorebestia* has jaws inside its head."

"This is what we see in microscopic jaw worms today—organisms that arrow worms shared an ancestor with over half a billion years ago. *Timorebestia* and other fossils like it provide links between closely related organisms that today look very different."

"Our discovery firms up how arrow worms evolved," added Tae Yoon Park from the Korean Polar Research Institute, the other senior author and field expedition leader. "Living arrow worms have a distinct nervous center on their belly, called a ventral ganglion. It is entirely unique to these animals."

"We have found this preserved in *Timorebestia* and another fossil called *Amiskwia*. People have debated whether or not *Amiskwia* was closely related to arrow [worms](#) as part of their evolutionary stem lineage. The preservation of these unique ventral ganglia gives us a great deal more confidence in this hypothesis."

"We are very excited to have discovered such unique predators in Sirius Passet. Over a series of expeditions to the very remote Sirius Passet in the furthest reaches of North Greenland, more than 82,5° north, we have collected a great diversity of exciting new organisms. Thanks to the remarkable, exceptional preservation in Sirius Passet, we can also reveal exciting anatomical details, including their digestive system, muscle anatomy, and nervous systems."

"We have many more exciting findings to share in the coming years that will help show how the earliest animal ecosystems looked like and evolved," Dr. Park concluded.

The work is [published](#) in the journal *Science Advances*.

**More information:** Tae-Yoon Park et al, A giant stem-group chaetognath, *Science Advances* (2024). [DOI: 10.1126/sciadv.adi6678](https://doi.org/10.1126/sciadv.adi6678).  
[www.science.org/doi/10.1126/sciadv.adi6678](https://www.science.org/doi/10.1126/sciadv.adi6678)

Provided by University of Bristol

Citation: 'Giant' predator worms more than half a billion years old discovered in North Greenland (2024, January 3) retrieved 2 May 2024 from <https://phys.org/news/2024-01-giant-predator-worms-billion-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.