


How HIV smuggles its genetic material into the cell nucleus

January 25 2024, by Carmen Rotte



Artist's impression shows how the HIV capsid penetrates the jelly-like permeability barrier of a nuclear pore. To smuggle its genome through this defense line into the cell nucleus, it has evolved into a molecular transporter. Credit: Johannes Pauly, Max Planck Institute for Multidisciplinary Sciences

Each year, about 1 million individuals worldwide become infected with

HIV, the virus that causes AIDS. To replicate and spread the infection, the virus must smuggle its genetic material into the cell nucleus and integrate it into a chromosome.

Research teams led by Dirk Görlich at the Max Planck Institute for Multidisciplinary Science and Thomas Schwartz at the Massachusetts Institute of Technology (MIT) have now discovered that its [capsid](#) has evolved into a molecular transporter. As such, it can directly breach a crucial barrier, which normally protects the [cell nucleus](#) against viral invaders. This way of smuggling keeps the [viral genome](#) invisible to anti-viral sensors in the cytoplasm. Their study is [published](#) in *Nature*.

Forty years after the [human immunodeficiency virus](#) (HIV) was discovered as the cause of AIDS, we have therapies that effectively keep the pathogen under control, but there is still no cure. The [virus](#) infects certain [immune cells](#) and hijacks their genetic program in order to multiply and replicate its own genetic material. The [infected cells](#) then produce the next generation of viruses until they are finally destroyed. The immunodeficiency symptoms of AIDS result from the massive loss of immune cells that normally fight viruses and other pathogens.

To use the host cell's resources, HIV must smuggle its genetic material through cellular defense lines into the cell nucleus. The nucleus, however, is closely guarded. Its [nuclear envelope](#) prevents unwanted proteins or harmful viruses from entering the nucleus and macromolecules from an uncontrolled escape. Yet, selected proteins can pass because the barrier is not hermetically sealed.

Thousands of tiny nuclear pores in the nuclear envelope provide a passageway. They control these transport processes with the help of importins and exportins—molecular transporters that capture cargoes with valid molecular "passcodes" and deliver them through the nuclear pore channel. A "smart" material turns these pores into one of nature's

most efficient sorting and transport machines.

'Smart' sorting in the nuclear pore

This "smart" material, called FG phase, is jelly-like and impenetrable for most macromolecules. It fills and blocks the nuclear pore channel. Importins and exportins, however, can pass through because their surfaces are optimized for sliding through an FG phase.

The cell's border control in the FG phase happens extremely fast—within milliseconds. Likewise, its transport capacity is enormous: A single nuclear pore can transfer up to 1,000 transporters per second through its channel. Even with such a high traffic density, the barrier of nuclear pores remains intact and keeps suppressing unwanted border crossings. HIV, however, subverts this control.

Smuggled genetic material

"HIV packages its genome into a capsid. Recent evidence suggests that the genome stays inside the capsid until it reaches the nucleus, and thus also when passing the nuclear pore. But there is a size problem," Thomas Schwartz of MIT explains. The central pore channel is 40 to 60 nanometers wide. The capsid has a width of about 60 nanometers and could just squeeze through the pore.

However, a normal cellular cargo would still be covered by a transporter layer that adds at least another ten nanometers. The HIV capsid would then be 70 nanometers wide—too big for a nuclear pore. "Nevertheless, [cryo-electron tomography](#) has shown that the HIV capsid gets into the nuclear pore. But how this happens has been so far a mystery in HIV infection," says Max Planck Director Görlich.

Camouflage as a molecular transporter

Together with Schwartz, Görlich has now discovered how the virus overcomes its size problem, namely by a sophisticated molecular adaptation. "The HIV capsid has evolved into a transporter with an importin-like surface. This way, it can slide through the FG phase of the nuclear pore. The HIV capsid can thus enter the nuclear pore without helping transporters and bypass the protective mechanism that otherwise prevents viruses from invading the cell nucleus," the biochemist explains.

His team has succeeded in reproducing FG phases in the laboratory. "Under the microscope, FG phases appear as micrometer-sized spheres that completely exclude normal proteins, but virtually suck up the HIV capsid with its enclosed contents," reports Liran Fu, one of the first authors of the study. "Similarly, the capsid is sucked up into the nuclear [pore](#) channel. This happens even after all cellular transporters have been removed."

In one respect the HIV capsid differs fundamentally from previously studied transporters that pass nuclear pores: It encapsulates its cargo completely and thus conceals its genomic payload from anti-viral sensors in the cytoplasm. Employing this trick, the viral [genetic material](#) can be smuggled through the cellular virus defense system without being recognized and destroyed. "This makes it another class of molecular transporters alongside importins and exportins," Görlich emphasizes.

There are still many unanswered questions, such as where and how the capsid disintegrates to release its contents. However, the observation that the capsid is an importin-like transporter might one day be exploited for better AIDS therapies.

More information: Liran Fu et al, HIV-1 capsids enter the FG phase

of nuclear pores like a transport receptor, *Nature* (2024). [DOI: 10.1038/s41586-023-06966-w](https://doi.org/10.1038/s41586-023-06966-w)

Provided by Max Planck Society

Citation: How HIV smuggles its genetic material into the cell nucleus (2024, January 25)
retrieved 27 April 2024 from
<https://phys.org/news/2024-01-hiv-smuggles-genetic-material-cell.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.