


# Methane pulses on Mars possibly driven by atmospheric pressure changes

January 24 2024

---



New simulations are helping inform the Curiosity rover's ongoing sampling campaign. Credit:NASA/JPL-Caltech/MSSS

New research shows that atmospheric pressure fluctuations that pull gases up from underground could be responsible for releasing subsurface

methane into Mars's atmosphere; knowing when and where to look for methane can help the Curiosity rover search for signs of life.

"Understanding Mars's [methane](#) variations has been highlighted by NASA's Curiosity team as the next key step towards figuring out where it comes from," said John Ortiz, a graduate student at Los Alamos National Laboratory who led the research team. "There are several challenges associated with meeting that goal, and a big one is knowing what time of a given sol (Martian day) is best for Curiosity to perform an atmospheric sampling experiment."

The paper was [published](#) Jan. 22 in the *Journal of Geophysical Research: Planets*.

A primary focus of NASA's Mars missions, including Curiosity and Perseverance, is to detect and understand past or present signs of life, such as methane. However, with the source of methane on Mars likely being underground, short-term variations in atmospheric methane levels have posed a research challenge.

To better understand Mars's methane levels, Ortiz and his team used high-performance computing clusters to simulate how methane travels through networks of underground fractures and is released into the atmosphere, where it then mixes within the atmospheric column. They also modeled how methane is adsorbed onto the pores of rocks, which is a temperature-dependent process that may contribute to the methane level fluctuations.

Their simulations predicted methane pulses from the [ground surface](#) into the atmosphere just before the Martian sunrise in the planet's northern summer season, which just recently ended. This corroborates previous rover data suggesting that methane levels fluctuated not only seasonally, but also daily.

This valuable data is helping inform the Curiosity rover's ongoing sampling campaign.

"Our work suggests several key time windows for Curiosity to collect data. We think these offer the best chance of constraining the timing of methane fluctuations, and (hopefully) down the line bringing us closer to understanding where it comes from on Mars," Ortiz said.

**More information:** J. P. Ortiz et al, Sub-Diurnal Methane Variations on Mars Driven by Barometric Pumping and Planetary Boundary Layer Evolution, *Journal of Geophysical Research: Planets* (2024). [DOI: 10.1029/2023JE008043](https://doi.org/10.1029/2023JE008043)

Provided by Los Alamos National Laboratory

Citation: Methane pulses on Mars possibly driven by atmospheric pressure changes (2024, January 24) retrieved 27 April 2024 from <https://phys.org/news/2024-01-methane-pulses-mars-possibly-driven.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.