


Molecular sensor enables water bear hardiness by triggering dormancy, study finds

January 17 2024



Tardigrade observed using a confocal fluorescent microscope. The tardigrade was overexposed to 5-MF, a cysteine selective fluorescent probe, that allows for visualization of internal organs. Credit: Smythers et al., 2024, PLOS ONE, CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)

Tardigrades—hardy, microscopic animals commonly known as "water bears"—use a molecular sensor that detects harmful conditions in their environment, telling them when to go dormant and when to resume normal life. A team led by Derrick R. J. Kolling of Marshall University and Leslie M. Hicks of the University of North Carolina at Chapel Hill report these findings in a study published in the open-access journal *PLOS ONE*.

Water bears are famous for their ability to withstand [extreme conditions](#), and can survive freezing, radiation, and environments without oxygen or water. They persist by going dormant and entering a tun state, in which their bodies become dehydrated, their eight legs retract and their metabolism slows to almost undetectable levels. Previously, little was known about what signals [water bears](#) to enter and leave this state.

In the new study, researchers exposed water bears to freezing temperatures or high levels of hydrogen peroxide, salt or sugar to trigger dormancy. In response to these harmful conditions, the animals' cells produced damaging oxygen free radicals.

The researchers found that water bears use a molecular sensor—based on the [amino acid cysteine](#)—which signals the animals to enter the tun state when it is oxidized by oxygen free radicals. Once conditions improve and the free radicals disappear, the sensor is no longer oxidized, and the water bears emerge from dormancy.

When the researchers applied chemicals that block cysteine, the water bears could not detect the [free radicals](#) and failed to go dormant.

Altogether, the new results indicate that cysteine is a key sensor for turning dormancy on and off in response to multiple stressors, including freezing temperatures, toxins and concentrated levels of salt or other compounds in the environment. The findings suggest that cysteine

oxidation is a vital regulatory mechanism that contributes to water bears' remarkable hardiness and helps them survive in ever-changing environments.

The authors add, "Our work reveals that tardigrade survival to stress conditions is dependent on reversible cysteine oxidation, through which [reactive oxygen species](#) serve as a sensor to enable tardigrades to respond to external changes."

More information: Chemobiosis reveals tardigrade tun formation is dependent on reversible cysteine oxidation, *PLoS ONE* (2024). [DOI: 10.1371/journal.pone.0295062](#)

Provided by Public Library of Science

Citation: Molecular sensor enables water bear hardiness by triggering dormancy, study finds (2024, January 17) retrieved 28 April 2024 from <https://phys.org/news/2024-01-molecular-sensor-enables-hardiness-triggering.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.