


Reaching the quantum ground state of sound in waveguides: Scientists move a step closer

January 18 2024





Artist's impression of cooled acoustic waves in an optical fiber taper. Credit: Long Huy Da

A team of scientists at the Max Planck Institute for the Science of Light led by Dr. Birgit Stiller has succeeded in cooling traveling sound waves in waveguides considerably further than has previously been possible using laser light. This achievement represents a significant move towards the ultimate goal of reaching the quantum ground state of sound in waveguides.

Unwanted noise generated by the acoustic waves at <u>room temperature</u> can be eliminated. This experimental approach both provides a deeper understanding of the transition from classical to quantum phenomena of <u>sound</u> and is relevant to quantum communication systems and future quantum technologies.

The quantum ground state of an acoustic wave of a certain frequency can be reached by completely cooling the system. In this way, the number of quantum particles, the so-called acoustic phonons, which cause disturbance to <u>quantum measurements</u>, can be reduced to almost zero and the gap between classical and <u>quantum mechanics</u> bridged.

Over the past decade, major technological advances have been made, making it possible to put various systems into this state. Mechanical vibrations oscillating between two mirrors in a resonator can be cooled to very low temperatures as far as the quantum ground state. This has not yet been possible for optical fibers in which high-frequency sound waves can propagate. Now, researchers from the Stiller Research Group have taken a step closer to this goal.

In their study, recently <u>published</u> in *Physical Review Letters*, they report




that they were able to lower the temperature of a <u>sound wave</u> in an <u>optical fiber</u> initially at room temperature by 219 K using laser cooling, ten times further than had previously been reported. Ultimately, the initial phonon number was reduced by 75% at a temperature of 74 K, -199 Celsius.

Such a drastic reduction in temperature was made possible by using laser light. Cooling of the propagating sound waves was achieved via the nonlinear optical effect of stimulated Brillouin scattering, in which <u>light</u> waves are efficiently coupled to sound waves.

Through this effect, the laser light cools the acoustic vibrations and creates an environment with less thermal noise, which is, to an extent, "disturbing" noise for a quantum communication system, for example. "An interesting advantage of glass fibers, in addition to this strong interaction, is the fact that they can conduct light and sound excellently over long distances," says Laura Blázquez Martínez, one of the lead authors of the article and a doctoral student in the Stiller research group.





Experimental setup in the laboratory. Credit: SAOT Max Gmelch

Most physical platforms previously brought to the quantum ground state were microscopic. However, in this experiment, the length of the optical fiber was 50 cm, and a sound wave extending over the full 50 cm of the core of the fiber was cooled to extremely low temperatures.

"These results are a very exciting step towards the quantum ground state in waveguides, and the manipulation of such long acoustic phonons opens up possibilities for broadband applications in quantum technology," according to Dr. Birgit Stiller, head of the quantum optoacoustics group.



Sound, in the day-to-day <u>classical world</u>, can be understood as a density wave in a medium. However, from the perspective of quantum mechanics, sound can also be described as a particle: the phonon. This particle, the sound quantum, represents the smallest amount of energy that occurs as an acoustic wave at a certain frequency. In order to see and study a single quanta of sound, the number of phonons must be minimized.

The transition from the classical to the quantum behavior of sound is often more easily observed in the quantum ground state, where the number of phonons is close to zero on average, such that the vibrations are almost frozen and quantum effects can be measured.

Stiller says, "This opens the door to a new landscape of experiments that allow us to gain deeper insights into the fundamental nature of matter." The advantage of using a waveguide system is that light and sound are not bound between two mirrors but propagating along the waveguide. The acoustic waves exist as a continuum—not only for specific frequencies—and can have a broad bandwidth, making them promising for applications such as high-speed communication systems.

"We are very enthusiastic about the new insights that pushing these fibers into the quantum ground state will bring," emphasizes the research group leader. "Not only from the fundamental research point of view, allowing us to peek into the quantum nature of extended objects, but also because of the applications this could have in quantum communications schemes and future quantum technologies."

More information: Laura Blázquez Martínez et al, Optoacoustic Cooling of Traveling Hypersound Waves, *Physical Review Letters* (2024). DOI: 10.1103/PhysRevLett.132.023603



Provided by Max Planck Institute for the Science of Light

Citation: Reaching the quantum ground state of sound in waveguides: Scientists move a step closer (2024, January 18) retrieved 9 May 2024 from https://phys.org/news/2024-01-quantum-ground-state-waveguides-scientists.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.