

New rice lines for Africa offer virus protection

January 11 2024, by Arne Claussen



Various rice plants, both control plants and mutants, between three and four weeks after inoculation. Credit: IRD / Laurence Albar

Rice yellow mottle virus (RYMV) is responsible for high crop losses in Africa, particularly among small-scale farmers. A research team has now

produced rice lines that are resistant to the disease by means of genome editing.

The [rice](#) varieties, the development of which the team describes in *Plant Biotechnology Journal*, are a preliminary step toward being able to generate resistant locally adapted elite varieties for small-scale food producers in Africa.

RYMV is an RNA virus spread by beetles and direct leaf-to-leaf contact. In Africa, where the majority of producers farm plots of land barely one hectare in size, between ten and one hundred percent of the rice harvests are regularly lost to this virus. This makes it a life-threatening problem for the poorest farmers.

There is no effective protection against the virus. "The only real protection is to develop rice varieties that possess a [resistance gene](#) against RYMV, which would make the plant invulnerable," says Dr. Yugander Arra, lead author of the study now [published](#) in *Plant Biotechnology Journal*.

A research team from the Institute for Molecular Physiology at HHU (headed by Professor Dr. Wolf B. Frommer) and the Institut de recherche pour le développement (IRD) in Montpellier, France, has developed such resistant rice lines.

Three [resistance genes](#) are currently known; mutations in just one gene, RYMV1, 2, and 3, are sufficient to achieve resistance. The resistant form *rymv2* occurs in poor-yielding African rice (*Oryza glaberrima*) varieties. RYMV2, also known as CPR5.1, encodes an essential protein from the pores of the cell nucleus.

In the model plant *Arabidopsis thaliana*, the loss of the only gene copy of CPR5 results in a broad spectrum of resistance to viruses, bacteria, and

fungi. However, growth is severely restricted; the plants exhibit spontaneous lesions and produce low yields. So, it was important to test whether rymv2 resistance could be transferred to other rice varieties without negative consequences.

In Africa, other high-yield rice varieties based on the Asian species *Oryza indica* are mainly used, and these do not have the resistance gene. Inserting the relevant gene is, however, not a particularly promising approach as the descendants of such "inter-species" hybrids are highly sterile and, therefore, cannot reproduce and pass on the resistance easily.

Using the CRISPR/Cas genome editing method, the research group has now shown that mutations of the RYMV2 gene can be produced in an Asian rice variety that make it resistant to the virus in a similar way to the African form. In the next step, the aim is to edit relevant African elite varieties in the same way to make them available to African small-scale producers. Helping these farmers is the goal of the international research consortium "Healthy Crops," which is headed by HHU.

Plants have hereditary mechanisms that were useful for survival in the early days of evolution but which are now more likely to be harmful. Maize is a good example of this: A gene causes the kernels to be aborted when drought conditions prevail at the time of fertilization. This trait caused by the gene was good for the wild perennial forebears of today's maize plants but has a [detrimental effect](#) on the yield of the annual plants now used in agriculture.

The situation is similar with the rice examined here. Professor Frommer says, "This resistance trait is attributable to the loss of a gene function that is not essential. If we switch the gene off completely, the plants behave normally. However, as a result of the loss of the gene function, they are resistant to the virus."

Dr. Eliza Loo, Healthy Crops Group Leader, adds, "It is so to speak an archetype, which was useful for its forebears, but which now leads to devastating crop losses in periods of drought. It would appear expedient to switch this gene off and it has no obvious side effects."

Surprisingly, neither switching off the closely related CPR5.2 gene nor the two [genes](#) RYMV 2 and CPR5.2—at least under greenhouse conditions—leads to impairments. It is also noteworthy that the loss of CPR5.2 does not lead to RYMV [resistance](#). Everything indicates that editing the RYMV2 gene is a promising approach to combating rice disease in Africa.

More information: Yugander Arra et al, Rice Yellow Mottle Virus resistance by genome editing of the *Oryza sativa* L. ssp. japonica nucleoporin gene OsCPR5.1 but not OsCPR5.2, *Plant Biotechnology Journal* (2023). [DOI: 10.1111/pbi.14266](https://doi.org/10.1111/pbi.14266)

Provided by Heinrich-Heine University Duesseldorf

Citation: New rice lines for Africa offer virus protection (2024, January 11) retrieved 14 May 2024 from <https://phys.org/news/2024-01-rice-lines-africa-virus.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--