


Unlocking the secrets of disease-causing fungus *Aspergillus fumigatus*

January 4 2024



GliT:GFP and GtmA:GFP have enriched vacuolar localization during GT production. GliT:GFP and GtmA:GFP germlings were grown in liquid Czapek-dox medium for 24 h at 37 °C. Representative brightfield, differential interference contrast (DIC), and Cell tracker Blue CMAC (CellTracker Blue CMAC Dye (7-amino-4-chloromethylcoumarin) for vacuolar staining. **a** The number of GliT:GFP and GtmA:GFP germlings that co-localized with CMAC that was used for vacuolar staining were determined. We have counted three independent experiments with 45 germlings for each strain per experiment ($N =$

135 germlings) and the results were expressed as the mean values (%) of 3 independent experiments of GFP that co-localizes with CMAC. Yellow arrows indicate the localization of GliT:GFP and GtmA:GFP in magnified images of sub-cellular structures. The p -values were calculated using One-way ANOVA with Tukey's multiple comparisons test, **** p

Citation: Unlocking the secrets of disease-causing fungus *Aspergillus fumigatus* (2024, January 4) retrieved 13 August 2024 from <https://phys.org/news/2024-01-secrets-disease-fungus-aspergillus-fumigatus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.