

## Single-emitter super-resolved imaging of radiative decay rate enhancement in dielectric gap nanoantennas

January 3 2024



smFLIM image of the decay rate enhancement induced by GaP dimers. Credit: *Light: Science & Applications* (2024). DOI: 10.1038/s41377-023-01349-2

In an era where understanding and manipulating light at the nanoscale is increasingly crucial, a paper in *Light: Science & Applications* reveals a significant leap forward.

A team of scientists from the Institut Langevin, ESPCI Paris, PSL University, CNRS have developed a sophisticated method to measure the enhancement of light interaction at the nanoscale using single molecules as probes. Central to this research are dielectric gap



nanoantennas—developed and fabricated at the Imperial College London.

Such structures are made of gallium phosphide (GaP), a material chosen for its high refractive index and low optical losses. This collaborative work involves an innovative approach using single molecules to probe the enhanced interaction of light facilitated purely by these nanoantennas without modification of the nanosystem with near-field probes, achieving a noticeable 30-fold enhancement in radiative decay rates at the single molecule level.

The scientists explain, "Our work focuses on the precise measurement of how light interacts with nanostructures. By using single <u>molecules</u> as probes, we've been able to observe and quantify the enhancement in light interaction, a crucial aspect for advancing nanophotonic technologies."

The research goes beyond mere theoretical exploration, offering practical insights into light-matter interactions. "This isn't just about observing enhanced light interaction; it's about measuring it at the singlemolecule level with remarkable spatial precision. Our findings are pivotal for future applications in fields where understanding and controlling light at such a small scale are essential."

The study's methodology and results underscore the effectiveness of advanced measurement techniques in nanophotonics.

"Our research has successfully mapped the spatial distribution of radiative decay rate enhancement, revealing that while there is some mislocalization of <u>single molecules</u> due to their interaction with the structure, this effect is minimal within the gap of the nanoantenna, providing a <u>precise control</u> of bright single-photon emission source," the scientists explain.



"This precision in measurement opens up new avenues for the characterization of highly sensitive optical devices and deepens our understanding of the interaction enhancement of a quantum emitter with a nanostructure."

In conclusion, the scientists emphasize the broader implications of their work. "Our research provides a new lens through which to view nanophotonic interactions. The ability to measure light interaction with such precision paves the way for breakthroughs in various applications, from <u>quantum computing</u>, quantum sensing to medical diagnostics."

**More information:** R. Margoth Córdova-Castro et al, Single-emitter super-resolved imaging of radiative decay rate enhancement in dielectric gap nanoantennas, *Light: Science & Applications* (2024). DOI: 10.1038/s41377-023-01349-2

Provided by Light Publishing Center, Changchun Institute of Optics, Fine Mechanics And Physics, CAS

Citation: Single-emitter super-resolved imaging of radiative decay rate enhancement in dielectric gap nanoantennas (2024, January 3) retrieved 21 May 2024 from <u>https://phys.org/news/2024-01-single-emitter-super-resolved-imaging-decay-dielectric.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.