

Team uses magnetized neurons to treat Parkinson's disease symptoms

January 18 2024



Credit: Nano Letters (2023). DOI: 10.1021/acs.nanolett.3c03899

Electrical deep brain stimulation (DBS) is a well-established method for treating disordered movement in Parkinson's disease. However, implanting electrodes in a person's brain is an invasive and imprecise way to stimulate nerve cells. Researchers <u>report</u> in *Nano Letters* on a new application for the technique, called magnetogenetics, that uses very small magnets to wirelessly trigger specific, gene-edited nerve cells in



the brain. The treatment effectively relieved motor symptoms in mice without damaging surrounding brain tissue.

In traditional DBS, a <u>battery pack</u> externally sends <u>electrical signals</u> through wires, activating <u>nerve cells</u> in a region of the brain called the subthalamic nucleus (STN). STN activation can relieve <u>motor symptoms</u> of Parkinson's disease, including tremors, slowness, rigidity and involuntary movements.

However, because the potential side effects, including brain hemorrhage and <u>tissue damage</u>, can be severe, DBS is usually reserved for people who have late-stage Parkinson's disease or when symptoms are no longer manageable with medication.

In a step toward a less invasive treatment, researchers Minsuk Kwak and Jinwoo Cheon worked with their colleagues to develop a wireless method to effectively reduce motor dysfunction in people with Parkinson's disease.

For their wireless technique, the researchers tagged nanoscale magnets with antibodies to help the molecules "stick" to the surface of STN nerve cells. Then they injected the sticky magnets into the brains of mice with early- and late-stage Parkinson's disease.

Prior to the injection in the STN, those same nerve cells had been modified with a gene that caused them to activate when the modified magnets on the cell's surface twisted in reaction to an externally applied magnetic field of about 25 milliteslas, which is about one- thousandth the strength of an MRI.

In demonstrations of the magnetized and modified neurons in mice with Parkinson's disease, the mice exposed to a magnetic field showed improved motor function to levels comparable to those of healthy mice.



The team observed that mice that received multiple exposures to the magnetic field retained more than one-third of their motor improvements while mice that received one exposure retained almost no improvements.

Additionally, the nerve cells of treated mice showed no significant damage in and around the STN, which suggests this could be a safer alternative to traditional implanted DBS systems, the researchers say. The team believes its wireless magnetogenetic approach has therapeutic potential and could be used to treat motor dysfunction in people with early- or late-stage Parkinson's disease as well as other neurological disorders, such as epilepsy and Alzheimer's disease.

More information: Wookjin Shin et al, Nanoscale Magnetomechanical-genetics of Deep Brain Neurons Reversing Motor Deficits in Parkinsonian Mice, *Nano Letters* (2023). DOI: <u>10.1021/acs.nanolett.3c03899</u>

Provided by American Chemical Society

Citation: Team uses magnetized neurons to treat Parkinson's disease symptoms (2024, January 18) retrieved 13 May 2024 from https://phys.org/news/2024-01-team-magnetized-neurons-parkinson-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.