

## Team shows how uranium ditelluride continues superconducting even in high magnetic fields

January 31 2024, by Simon Schmitt



Graphic representation of a microstructured sample (red) for electrical measurements on unconventional superconductors. Gold and platinum are used for contacting. Electrons (green spheres) couple in pairs via vibrational or magnetic fluctuations. Credit: B. Schröder/ HZDR

Superconductivity is well understood in so-called "conventional" superconductors. More recent, however, are unconventional



superconductors, and it is as yet unclear how they work.

A team from HZDR, together with colleagues from CEA, the Tohoku University in Japan, and the Max Planck Institute for Chemical Physics of Solids, has now explained why a <u>new material</u> continues superconducting even in extremely high magnetic fields—a property that is missing in conventional superconductors. The finding has the potential to enable previously inconceivable technological applications. The study is <u>published</u> in *Nature Communications*.

"Uranium ditelluride, or UTe<sub>2</sub> for short, is a high-flyer among <u>superconducting materials</u>," says Dr. Toni Helm from the Dresden High Magnetic Field Laboratory (HLD) at HZDR. "As discovered in 2019, the compound conducts electricity without loss, however, in a different way than conventional superconductors do."

Since then, research groups around the world have become interested in the material. This includes Helm's team, which has come a step closer to understanding the compound.

"To fully appreciate the hype surrounding the material, we need to take a closer look at superconductivity," explains the physicist. "This phenomenon results from the movement of electrons in the material. Whenever they collide with atoms, they lose energy in form of heat. This manifests itself as electrical resistance. Electrons can avoid this by arranging themselves in pair formations, so-called Cooper pairs."

A Cooper pair describes two electrons combined at <u>low temperatures</u> to move through a solid without friction. They make use of the atomic vibrations around them as a kind of wave on which they can surf without losing energy. These atomic vibrations explain conventional superconductivity.



"For some years now, however, superconductors have also been known in which Cooper pairs are formed by effects that are not yet fully understood," says the physicist. One possible form of unconventional superconductivity is spin-triplet superconductivity, which is believed to make use of magnetic fluctuations.

"There are also metals in which the conduction electrons come together collectively," explains Helm. "Together, they can shield the magnetism of the material, behaving as a single particle with—for electrons—an extremely high mass."

Such superconducting materials are known as heavy-fermion superconductors.  $UTe_2$ , therefore, could be both a spin-triplet and a heavy-fermion superconductor, as current experiments suggest. Beyond this, it is the heavyweight world champion—to date, no other known heavy-fermion superconducts at similar or higher magnetic fields. This too was confirmed by the present study.

## Extremely robust against magnetic fields

Superconductivity depends on two factors: The critical transition temperature and the critical magnetic field. If the temperature falls below the critical transition temperature, the resistance drops to zero and the material becomes superconducting. External magnetic fields also influence superconductivity. If these exceed a critical value, the effect collapses.

"Physicists have a rule of thumb for this," said Helm. "In many conventional superconductors, the value of the transition temperature in Kelvin is roughly one to two times the value of the critical magneticfield strength in tesla. In spin-triplet superconductors, this ratio is often much higher."



With their studies on the heavyweight  $UTe_2$ , the researchers have now been able to raise the bar even higher: At a transition temperature of 1.6 kelvin (-271.55°C), the critical magnetic-field strength reaches 73 tesla, setting the ratio at 45—which is a record.

"Until now, heavy-fermion superconductors were of little interest for technical applications," explains the physicist. "They have a very low transition temperature and the effort required to cool them is comparatively high."

Nevertheless, their insensitivity to external magnetic fields could compensate for this shortcoming. This is because lossless current transport is mainly used today in <u>superconducting magnets</u>, for example in magnetic-resonance-imaging (MRI) scanners. However, the magnetic fields also influence the superconductor itself.

A material that can withstand very high magnetic fields and still conduct electricity without loss would represent a major step forward.

## Special treatment for a demanding material

"Of course, UTe<sub>2</sub> cannot be used to make leads for a superconducting electromagnet," says Helm. "First, the material's properties make it unsuitable for this endeavor, and second, it is radioactive. But it is perfectly suited for the exploration of the physics behind spin-triplet superconductivity."

Based on their experiments, the researchers developed a model that could serve as an explanation for superconductivity with extremely high stability against magnetic fields. To do this, they worked on samples with thicknesses of a few micrometers—only a fraction of the thickness of a human hair (approximately 70 micrometers). The radioactive radiation emitted by the samples, therefore, remains much lower than



that of the natural background.

In order to obtain and shape such a tiny sample, Helm used a highprecision ion beam with a diameter of just a few nanometers as a cutting tool.  $UTe_2$  is an air-sensitive material. Consequently, Helm carries out the sample preparation in a vacuum and seals them in epoxide glue afterwards.

"For the final proof that our material is a spin-triplet superconductor, we would have to examine it spectroscopically while it is exposed to strong magnetic fields. However, current spectroscopy methods still struggle at magnetic fields above 40 tesla. Alongside other teams, we are also working on developing novel techniques. Eventually, this will enable us to provide definitive proof," says Helm.

**More information:** Toni Helm et al, Field-induced compensation of magnetic exchange as the possible origin of reentrant superconductivity in UTe<sub>2</sub>, *Nature Communications* (2024). DOI: 10.1038/s41467-023-44183-1

## Provided by Helmholtz Association of German Research Centres

Citation: Team shows how uranium ditelluride continues superconducting even in high magnetic fields (2024, January 31) retrieved 8 May 2024 from <u>https://phys.org/news/2024-01-team-uranium-ditelluride-superconducting-high.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.