

From New York to Jakarta, land in many coastal cities is sinking faster than sea levels are rising

January 25 2024, by Pei-Chin Wu, Meng (Matt) Wei and Steven D'Hondt



Credit: Pixabay/CC0 Public Domain

<u>Sea level rise</u> has already put coastal cities on notice thanks to increasing storm surges and even <u>sunny day</u> flooding at high tide. These challenges



will continue to grow because global projections point to a mean sea level rise of at least one foot above year-2000 levels by the <u>end of this</u> <u>century</u>.

However, many cities are facing another factor making them even more vulnerable to rising waters: <u>land subsidence</u>.

The three of us—<u>Pei-Chin Wu</u>, <u>Meng (Matt) Wei</u> and <u>Steven D'Hondt</u> —are scientists at the University of Rhode Island Graduate School of Oceanography working with the U.S. Geological Survey to research challenges facing waterfront cities. Our findings indicate that <u>land is</u> <u>sinking</u> faster than sea levels are rising in many coastal cities throughout the world.

By using radar images of the Earth's surface collected from orbiting satellites, we measured subsidence rates in 99 <u>coastal cities</u> worldwide. These rates are highly variable within cities and from city to city, but if they continue, many metropolises will experience flooding much sooner than projected by <u>sea level rise</u> models.

Cities in South, Southeast and East Asia are seeing the most rapid rates of subsidence.

Indonesia, for example, is moving its capital 800 miles from Jakarta to Nusantara in large part because Jakarta is sinking at an alarming rate due to groundwater extraction.

Other regions are not immune. Our research with Tom Parsons of the U.S. Geological Survey found that most of <u>New York City</u>is sinking between 1 to 4 millimeters per year due to a combination of <u>glacial</u> rebound and the weight of its more than 1 million buildings. In a city where sea level is projected to rise between 8 and 30 inches by 2050, subsidence <u>further increases its vulnerability</u> to coastal storms.



In the U.S., most of the cities on the Atlantic coast are subsiding due to glacial rebound. Even if the rate is low at minus-1 millimeter per year, it should be accounted. Other cities in the U.S., especially in the Gulf of Mexico, including Houston and New Orleans, also face subsidence.

Governments around the world are facing the challenge of coastal areas that are subsiding, and there is a shared global challenge of mitigation against a growing flooding hazard.

While our research continues to evolve—for example, by using machine learning to improve our monitoring capability—we urge <u>city planners</u>, emergency managers and other <u>decision-makers</u> to account for subsidence in the plans they are making today to prepare for the impacts of rising sea levels in the future.

Correction note: This story was updated to correct the rate at which New York City is sinking.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: From New York to Jakarta, land in many coastal cities is sinking faster than sea levels are rising (2024, January 25) retrieved 12 May 2024 from <u>https://phys.org/news/2024-01-york-jakarta-coastal-cities-faster.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.