

Carbon emissions from the destruction of mangrove forests predicted to increase by 50,000% by the end of the century

February 22 2024



Credit: Unsplash/CC0 Public Domain

The annual rate of carbon emissions due to the degradation of carbon stocks in mangrove forests is predicted to rise by nearly 50,000% by the



end of the century, according to a new study <u>published</u> in *Environmental Research Letters*. Mangroves in regions such as southern India, southeastern China, Singapore and eastern Australia are particularly affected.

Mangrove forests store a large amount of carbon, particularly in their soils, however human development in these areas has led to the degradation of these carbon stocks. Over the past 20 years, a substantial number of <u>mangrove forests</u> have been replaced by agriculture, aquaculture and urban land management, leading global mangrove carbon stocks to decline by 158.4 million tons—releasing the same level of carbon emissions as flying the entire US population from New York to London.

The study is led by Jennifer Krumins, Professor of Biology at Montclair State University, alongside two Ph.D. candidates, Shih-Chieh Chien and Charles Knoble. The team focused on the relationship between <u>human</u> <u>population density</u> and soil carbon stocks in urban mangrove forests to quantify their role in the global carbon budget.

The results show that when population density reaches 300 people/km² (similar to the average population density of the UK or Japan), the carbon stored in mangrove soils near populated areas is estimated to be 37% lower than isolated mangrove forests. At the same time, the annual rate of carbon emissions from mangrove loss is currently estimated to be 7.0 Teragrams, rising in line with increased population density to 3,392 Teragrams by current predictions at the end of the century.

Mangrove forests cover approximately 0.1% of the Earth's land surface but play a vital role in providing wildlife habitats and regulating global climate stability. These mangroves store a large amount of carbon, particularly in their soils, and are essential to regulating carbon cycling on a global scale. Mangrove soils contain three to four times the mass of



carbon typically found in boreal, temperate, or tropical forests.

Professor Krumins says, "This work underscores the importance of protecting existing mangroves, especially in areas with high population density. Mangrove forests are critical to the regulation of carbon sequestration, and it is important that we protect them. The first step is to understand the impact of human populations and activities on mangrove forest carbon stocks."

More information: *Environmental Research Letters* (2024). DOI: 10.1088/1748-9326/ad13b6. iopscience.iop.org/article/10. 088/1748-9326/ad13b6

Provided by IOP Publishing

Citation: Carbon emissions from the destruction of mangrove forests predicted to increase by 50,000% by the end of the century (2024, February 22) retrieved 8 May 2024 from https://phys.org/news/2024-02-carbon-emissions-destruction-mangrove-forests.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.