


# Discovery of primitive mitochondrial DNA replication enzymes

February 28 2024



Subcellular localizations of GFP fused with the N-termini of 5 rdxPolAs when expressed in yeast cells. GFP fused with the N-terminal amino acid residues

predicted as the MTS of rdxPolA proteins from 3 members of Discoba (Ophirina amphinema, Tsukubamonas globosa, and Naegleria gruberi; a to c), the malawimonad (Gefionella okellyi; d), and the ancyromonad (Fabomonas tropica; e). The green signal corresponds to GFP. The red color indicates mitochondria stained by MitoTracker Red. The results presented here demonstrate that the mitochondrial translocons in yeast recognize the N-termini of the 5 rdxPolAs as the MTS. Credit: *Molecular Biology and Evolution* (2024). DOI: 10.1093/molbev/msae014

Researchers led by University of Tsukuba have discovered rdxPolA, a putative DNA polymerase involved in replicating ancestral mitochondrial genomes, in diverse eukaryotic lineages. Based on the phylogenetic distribution of rdxPolA among eukaryotes, they proposed an evolutionary scenario of DNA polymerases for mitochondrial genome maintenance in the early evolution of eukaryotes.

Mitochondria are intracellular organelles that evolved from a [bacterium](#) belonging to Alphaproteobacteria, which was taken up as an endosymbiont by the common ancestor of eukaryotes. Mitochondria possess their own highly reduced genomes (known as mitochondrial genomes), which are principally the descendants of the genome of the  $\alpha$ -proteobacterial symbiont. Phylogenetically diverse eukaryotes use a type of DNA polymerase called "POP" to maintain their mitochondrial genomes.

In this study, [published](#) in *Molecular Biology and Evolution*, the researchers identified 10 novel types of DNA polymerase that are distinct from the previously known types, including POPs, across diverse eukaryotic lineages. The [evolutionary origin](#) and subcellular localization of each novel DNA polymerase were investigated.

Intriguingly, one of the DNA polymerases identified in this study,

rdxPolA, was found to be involved in mitochondrial DNA maintenance and is a direct descendant of the DNA polymerase in the  $\alpha$ -proteobacterial symbiont that gave rise to the first mitochondrion.

The researchers proposed a scenario for the evolution of DNA polymerases involved in mitochondrial DNA maintenance from primitive to extant eukaryotes.

These findings provide critical insights into the early evolution of the machinery for mitochondrial DNA maintenance and the establishment of [mitochondria](#) in primitive eukaryotic cells.

**More information:** Ryo Harada et al, Encyclopedia of Family A DNA Polymerases Localized in Organelles: Evolutionary Contribution of Bacteria Including the Proto-Mitochondrion, *Molecular Biology and Evolution* (2024). [DOI: 10.1093/molbev/msae014](https://doi.org/10.1093/molbev/msae014)

Provided by University of Tsukuba

Citation: Discovery of primitive mitochondrial DNA replication enzymes (2024, February 28) retrieved 27 April 2024 from

<https://phys.org/news/2024-02-discovery-primitive-mitochondrial-dna-replication.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--