


Research confirms exogenous methyl jasmonate can enhance tomato resistance

February 27 2024




Binding of SIWRKY80 to the promoter region of *SISPDS2* and promotion of Spd synthesis. Credit: *Horticulture Research*

Tomato (*Solanum lycopersicum* L.) is the most widely cultivated and consumed horticultural crop. At present, saline-alkali is an important abiotic stress source that affects tomato production. Exogenous methyl jasmonate (MeJA) can enhance the resistance of tomatoes to various stresses, but its exact mechanism is still unclear.

Horticulture Research has now [published](#) new research titled "SIWRKY80-mediated jasmonic acid pathway positively regulates tomato resistance to saline-alkali stress by enhancing spermidine content and stabilizing Na⁺/K⁺ homeostasis."

In this study, researchers confirmed that 22.5 µmol/l MeJA could significantly improve the saline-alkali stress resistance of tomatoes. Saline-alkali stress increased the endogenous MeJA and [jasmonic acid](#) (JA) contents. Exogenous application of 22.5 µmol/l MeJA increased the endogenous MeJA and JA contents in tomato.

Furthermore, an important transcription factor, SIWRKY80, responded to MeJA and actively regulated tomato resistance to saline-alkali stress. Spraying of exogenous MeJA (22.5 µmol/l) reduced the sensitivity of SIWRKY80 knockout lines to saline-alkali stress.



A working model for the saline-alkali reaction mediated by SIWRKY80 through the JA pathway and the regulation of *SISPDS2* and *SINHX4* in tomato. Credit: *Horticulture Research*

The SIWRKY80 protein directly combines with the promoter of *SISPDS2* and *SINHX4* to positively regulate the transcription of *SISPDS2* and *SINHX4*, thereby promoting the synthesis of spermidine and Na^+/K^+ homeostasis, actively regulating saline-alkali stress. The augmentation of JA content led to a notable reduction of 70.6% in the expression of SIJAZ1 and the release of the SIWRKY80 protein interacting with SIJAZ1.

In conclusion, exogenous MeJA in tomato stress resistance through multiple [metabolic pathways](#) elucidated that exogenous MeJA further promotes spermidine synthesis and Na^+/K^+ [homeostasis](#) by activating the expression of SIWRKY80, which provides a new theoretical basis for the study of the JA stress [resistance](#) mechanism and the actual production of tomato.

More information: Chunyu Shang et al, SIWRKY80-mediated JA pathway positively regulates tomato resistance to saline-alkali stress by enhancing spermidine content and stabilizing Na⁺/K⁺ homeostasis, *Horticulture Research* (2024). [DOI: 10.1093/hr/uhae028](https://doi.org/10.1093/hr/uhae028)

Provided by NanJing Agricultural University

Citation: Research confirms exogenous methyl jasmonate can enhance tomato resistance (2024, February 27) retrieved 2 May 2024 from <https://phys.org/news/2024-02-exogenous-methyl-jasmonate-tomato-resistance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.