


Impact of moisture sources on variability of precipitation oxygen isotopes in northern China

February 19 2024



Spatial correlation fields of CAM3 simulated $\delta^{18}\text{O}$ in northern China and outgoing longwave radiation. **a** Spatial correlations between CAM3 simulated JJA amount weighted precipitation $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_p$) in northern China and JJA outgoing longwave radiation (OLR) at each grid cell. **b** Same as **a** but for CAM3 simulated water vapor $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_v$) at lower-troposphere (1000–750 hPa) in JJA. Credit: *npj Climate and Atmospheric Science* (2024). DOI: 10.1038/s41612-024-00564-x

In a [study](#) published in *npj Climate and Atmospheric Science*, a joint research group from the Institute of Earth Environment of the Chinese

Academy of Sciences, the Stockholm University, and the California State University found that while land recycled water vapor contributes more than 60% to precipitation in northern China, the precipitation $\delta^{18}\text{O}$ ($\delta^{18}\text{Op}$) is primarily influenced by the relative contribution of oceanic moisture.

For decades, precipitation $\delta^{18}\text{O}$ ($\delta^{18}\text{Op}$) has been an important proxy for reconstructing past climate change. However, its interpretation, especially under the influence of the East Asian Summer Monsoon (EASM), has been controversial. In northern China, located at the margin of the EASM region, the more continental climate raises questions about the impact of different moisture sources on $\delta^{18}\text{Op}$.

To explore the complexity of moisture source variability and its impact on $\delta^{18}\text{Op}$, the researchers used a general circulation model the Community Atmosphere Model version 3 (CAM3), enhanced with water-tagging and stable water isotope modules, to quantitatively trace water vapor from its source to northern China.

They found that more than 60% of the precipitation in northern China originated from local recycled moisture and other terrestrial [water vapor](#). However, this substantial contribution had a minimal effect on $\delta^{18}\text{Op}$ values, indicating a predominant transpiration over evaporation.

They suggested that despite the high percentage of terrestrial moisture, the $\delta^{18}\text{Op}$ was predominantly affected by oceanic moisture sources, with the Pacific Ocean (24.4%) and the North Indian Ocean (13.0%) being crucial contributors.

In addition, the researchers highlighted the role of the Pacific Decadal Oscillation in modulating $\delta^{18}\text{Op}$ by altering the moisture contributions of the Pacific Ocean and North Indian Ocean.

Interestingly, local [precipitation](#) patterns show an inverse phase variation with $\delta^{18}\text{O}_p$, highlighting the complex dynamics between local moisture recycling and the large-scale oceanic influences.

These findings are valuable for understanding past climate variability and provide a new lens through which to view the complex climate systems of monsoonal China.

More information: Fangyuan Lin et al, Seasonal to decadal variations of precipitation oxygen isotopes in northern China linked to the moisture source, *npj Climate and Atmospheric Science* (2024). [DOI: 10.1038/s41612-024-00564-x](#)

Provided by Chinese Academy of Sciences

Citation: Impact of moisture sources on variability of precipitation oxygen isotopes in northern China (2024, February 19) retrieved 9 May 2024 from <https://phys.org/news/2024-02-impact-moisture-sources-variability-precipitation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--