


Exploring exotic behaviors in population-imbalanced fermionic systems

March 12 2024, by Samuel Jarman



Credit: *The European Physical Journal B* (2024). DOI: 10.1140/epjb/s10051-024-00649-9

Over the past 20 years, many physicists have studied ultra-cold fermionic systems contained in magnetic or optical traps. When an external magnetic field is applied to a two-species fermionic system, the particles can pair together to form composite "bosonic molecules" with a full-integer spin.

These molecules undergo Bose-Einstein condensation when cooled, where all the particles accumulate in the lowest-energy quantum state. The precision of these experiments has now been improved by trapping the particles inside optical lattices: periodic patterns formed by counter-propagating [laser beams](#).

Through research [published](#) in *The European Physical Journal B*, Avinaba Mukherjee and Raka Dasgupta at the University of Calcutta, India, have theoretically predicted a distinctive trend in the oscillations of Bose-Einstein condensates formed from these fermions, which can be adjusted using an [external magnetic field](#).

They specifically addressed systems where the two species have unequal populations (creating leftover unpaired fermions), leading to exotic new phases. Their result could help physicists to detect such novel phases of matter in imbalanced fermionic systems and could open up new opportunities for quantum technologies.

In their work, Mukherjee and Dasgupta explored how such a system would behave when they applied a technique often used to manipulate and control ultracold atomic gases. Named Feshbach detuning, it involves adjusting the energy required for forming bosonic molecules, using an external magnetic field.

The researchers discovered that when Feshbach detuning is above a certain threshold, the fraction of Bose-condensed particles will oscillate periodically—but below this threshold, there is no oscillation at all.

Altogether, this revealed a linear relationship between the oscillation frequency, and the strength of Feshbach detuning they applied.

In addition, the duo found that the slope and position of this curve carried important information about which exotic phase of matter could be found in the system. Their result may eventually lead to discoveries of advanced new physical properties in these systems, which could be exploited across a diverse range of quantum technologies.

More information: Avinaba Mukherjee et al, Periodic dynamics of population-imbalanced fermionic condensates in optical lattices, *The European Physical Journal B* (2024). [DOI: 10.1140/epjb/s10051-024-00649-9](https://doi.org/10.1140/epjb/s10051-024-00649-9)

Provided by SciencePOD

Citation: Exploring exotic behaviors in population-imbalanced fermionic systems (2024, March 12) retrieved 16 June 2024 from <https://phys.org/news/2024-03-exploring-exotic-behaviors-population-imbalanced.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.