


Frequency of heat days may be systematically underestimated in many studies

March 19 2024, by Theresa Bittermann



The study shows that heat days are underestimated by up to 30 percent on average over 30 years if the error is not corrected—in other words, only 70 percent of the actual temperature extremes are detected in some regions. Credit: Lukas Brunner

Many studies on the climate crisis focus on researching temperature extremes on a global scale. Scientists at the University of Vienna have now uncovered an error in an established calculation method, leading to a systematic underestimation in the frequency of heat days.


The error is based on the previously unnoticed impact of the seasonal cycle on the extreme threshold due to the incorrect application of so-called "moving time windows." The study has recently been [published](#) in the journal *Nature Communications*.

Increasingly frequent temperature extremes are one of the most dangerous consequences of human-induced climate change and, as such, the subject of numerous scientific analyses.

A commonly used method to define extremes such as heat days takes the adaptation to local conditions into account and calculates them relative to the local temperature distribution. However, researchers from the Department of Meteorology and Geophysics at the University of Vienna have now identified a significant error in the calculation of such relative extremes.

Temperature extremes are often defined relative to local conditions to encompass unusually hot periods worldwide. This approach uses different threshold values, for example, for Europe and Antarctica, allowing for a comparison in the occurrence of heat days between these climatically diverse regions. So-called moving time windows are often employed to calculate the local temperature threshold.

These windows aim to increase the number of days considered for threshold calculation, intending to enhance the meaningfulness of the threshold. Many previous studies have, therefore, increased the length of this time window from the originally recommended 5 days to up to 31 days. The newly published study now demonstrates that such long time windows lead to a mixing of the seasonal cycle into the threshold, inadvertently reducing the probability of extremes.



The impact of the error varies greatly from region to region—the effects are particularly clear in the North Atlantic: In June, for example, the frequency of hot days was underestimated by more than 75 percent, resulting in only a quarter of the actual extremes being detected. Averaged over the year, the temperature extremes were also significantly below the expected frequency. Credit: Lukas Brunner

Frequency of heat days underestimated

The calculation error can, depending on the region, lead to an underestimation of the expected heat day frequency, as explained by the lead author of the study, Lukas Brunner, Senior Scientist at the Department of Meteorology and Geophysics at the University of Vienna, "Heat days are often defined as the 10 percent warmest days at each location."

"However, we were able to show that an error in the calculation can lead to a considerable underestimation in the number of extreme days. This

has been overlooked in many studies so far."

Regions particularly affected include the western United States and the Arabian Peninsula, where only 7 percent heat days are detected by the algorithm instead of the correct 10 percent, leading to a relative error of -30 percent. In contrast, the probability of heat days in Europe is quite accurately estimated at the correct 10 percent.

"These [regional differences](#) in the manifestation of the error we have identified can distort the interpretation of results and lead to issues when comparing different regions of the world," explains Brunner.

Global warming reduces the error

The study also examines the [impact of climate change](#), revealing that with unabated emissions, certain regions will experience almost continuous temperature extremes in the future. The discovered underestimation of extremes decreases with extreme climate warming.

However, this leads to an overestimation of the change signal; as climate scientist Brunner explains, "By the end of the century, in hotspot regions like the Arabian Peninsula, almost every day will be considered a heat day by today's standards. But due to the error, the historical period has only 7 percent heat days instead of the correct 10 percent, leading to an overestimation in the increase."

In their study, the authors also propose a correction that almost entirely eliminates the error. "We hope that our study leads to future work avoiding the error, allowing for a better characterization of changes in [temperature extremes](#) within the framework of climate change," says Aiko Voigt, co-author of the study and Professor at the Department of Meteorology and Geophysics at the University of Vienna.

More information: Lukas Brunner et al, Pitfalls in diagnosing temperature extremes, *Nature Communications* (2024). [DOI: 10.1038/s41467-024-46349-x](https://doi.org/10.1038/s41467-024-46349-x)

Provided by University of Vienna

Citation: Frequency of heat days may be systematically underestimated in many studies (2024, March 19) retrieved 22 June 2024 from <https://phys.org/news/2024-03-frequency-days-systematically-underestimated.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.