

A new ion trap for larger quantum computers

March 13 2024, by Oliver Morsch



The experimental setup of the ETH researchers. The trap chip is located inside the container underneath the silver cupola, in which a lens captures the light emitted by the trapped ions. Credit: ETH Zurich / Pavel Hrmo

Researchers at ETH have managed to trap ions using static electric and magnetic fields and to perform quantum operations on them. In the future, such traps could be used to realize quantum computers with far more quantum bits than have been possible up to now.



The energy states of electrons in an atom follow the laws of quantum mechanics: They are not continuously distributed but restricted to certain well-defined values—this is also called quantization. Such quantized states are the basis for quantum bits (qubits), with which scientists want to build extremely powerful quantum computers. To that end, the atoms have to be cooled down and trapped in one place.

Strong trapping can be achieved by ionizing the atoms, which means giving them an <u>electric charge</u>. However, a fundamental law of electromagnetism states that electric fields that are constant in time cannot trap a single charged particle. By adding an oscillating electromagnetic field, on the other hand, one obtains a stable ion trap, also known as a Paul trap.

In this way, it has been possible in recent years to build quantum computers with ion traps containing around 30 qubits. Much larger quantum computers, however, cannot straightforwardly be realized with this technique. The oscillating fields make it difficult to combine several such traps on a single chip, and using them heats up the trap—a more significant problem as systems get larger. Meanwhile transport of ions is restricted to pass along linear sections connected by crosses.





Schematic showing the middle section of the used Penning trap. An ion (red) is trapped through a combination of an electric field produced by different electrodes (yellow) and a magnetic field. Credit: ETH Zürich / Institute for Quantum Electronics

Ion trap with a magnetic field

A team of researchers at ETH Zurich led by Jonathan Home has now demonstrated that ion traps suitable for use in quantum computers can also be built using static magnetic fields instead of oscillating fields. In those static traps with an additional magnetic field, called Penning traps, both arbitrary transport and the necessary operations for the future super-



computers were realized. The researchers recently <u>published</u> their results in the scientific journal *Nature*.

"Traditionally, Penning traps are used when one wants to trap very many ions for precision experiments, but without having to control them individually," says Ph.D. student Shreyans Jain. "By contrast, in the smaller quantum computers based on ions, Paul traps are used."

The idea of the ETH researchers to build future quantum computers also using Penning traps was initially met with skepticism by their colleagues for various reasons. Penning traps require extremely strong magnets, which are very expensive and rather bulky.

Also, all previous realizations of Penning traps had been very symmetric, something that the chip-scale structures used at ETH violate. Putting the experiment inside a large magnet makes it difficult to guide the <u>laser</u> <u>beams</u> necessary for controlling the qubits into the trap, while strong magnetic fields increase the spacing between the energy states of the qubits. This, in turn, makes the control laser systems much more complex: instead of a simple diode laser, several phase-locked lasers are needed.





Moving a single trapped ion in a two-dimensional plane and illuminating it with a laser beam allows the researchers to create the ETH logo. The image is formed averaging over many repetitions of the transport sequence. Credit: ETH Zurich / Institute for Quantum Electronics

Transport in arbitrary directions

Home and his collaborators were not deterred by those difficulties, however, and constructed a Penning trap based on a superconducting



magnet and a microfabricated chip with several electrodes, which was produced at the Physikalisch-Technische Bundesanstalt in Braunschweig. The magnet used delivers a field of 3 Tesla, almost 100,000 times stronger than Earth's magnetic field. Using a system of cryogenically cooled mirrors, the Zurich researchers managed to channel the necessary laser light through the magnet to the ions.

The efforts paid off: A single trapped ion, which can stay in the trap for several days, could now be moved arbitrarily on the chip, connecting points "as the crow flies" by controlling the different electrodes—this was not previously possible with the old approach based on oscillating fields. Since no oscillating fields are needed for trapping, many of those traps can be packed onto a single chip.

"Once they are charged up, we can even completely isolate the electrodes from the outside world and thus investigate how strongly the ions are disturbed by external influences," says Tobias Sägesser, who was involved in the experiment as a Ph.D. student.

Coherent control of the qubit

The researchers also demonstrated that the qubit energy states of the trapped ion could also be controlled while maintaining quantum mechanical superpositions. Coherent control worked both with the electronic (internal) states of the ion and the (external) quantized oscillation states as well as for coupling the internal and external quantum states. This latter is a prerequisite for creating entangled states, which are important for quantum computers.

As a next step, Home wants to trap two ions in neighboring Penning traps on the same chip and thus demonstrate that <u>quantum operations</u> with several qubits can also be performed. This would be the definitive proof that quantum computers can be realized using ions in Penning



traps. The professor also has other applications in mind. For instance, since the ions in the new trap can be moved flexibly, they can be used to probe electric, magnetic or microwave fields near surfaces. This opens up the possibility to use these systems as atomic sensors of surface properties.

More information: Shreyans Jain et al, Penning micro-trap for quantum computing, *Nature* (2024). DOI: 10.1038/s41586-024-07111-x

Provided by ETH Zurich

Citation: A new ion trap for larger quantum computers (2024, March 13) retrieved 23 June 2024 from <u>https://phys.org/news/2024-03-ion-larger-quantum.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.