


LHCb observes a new decay mode of the charmed beauty meson

March 4 2024



The LHCb detector. Credit: Maximilien Brice, CERN

The LHCb collaboration recently [reported](#) on the *arXiv* preprint server the first observation of the decay of the B_c^+ meson (composed of two heavy quarks, b and c) into a J/ψ charm-anticharm quark bound state

and a pair of pions, $\pi^+\pi^0$. The decay process shows a contribution from an intermediate particle, a ρ^+ meson that forms for a brief moment and then decays into the $\pi^+\pi^0$ pair.

The B_c^+ is the heaviest meson that can only decay through the [weak interactions](#), via the decay of one heavy constituent quark. B_c^+ decays into an odd number of light hadrons and a J/ψ (or other charm-anticharm quark bound states, called "charmonia") have been studied intensively and have been found to be in remarkable agreement with the theoretical expectations.

The decay of B_c^+ into a J/ψ and a $\pi^+\pi^0$ pair is the simplest decay into charmonium and an even number of light hadrons. It has never been observed before, mainly because the precise reconstruction of the low-energy π^0 meson through its decay into a pair of photons is very challenging in an LHC proton-proton collision environment.

A precise measurement of the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decay will allow better understanding of its possible contribution as a background source for the study of other decays of B_c mesons as well as rare decays of B^0 mesons. From the theoretical point of view, decays of B_c into J/ψ and an even number of pions are closely related to the decays of the τ lepton into an even number of pions, and to the e^+e^- [annihilation](#) into an even number of pions.

Precise measurements of e^+e^- annihilation into two pions in the ρ mass region (as in the B_c decay discussed here) are crucial for the interpretation of results from the Fermilab $g-2$ experiment measuring the anomalous magnetic dipole moment of the muon, since low-energy e^+e^- annihilation into hadrons is an important source of the uncertainty of the $g-2$ measurements.

The ratio of the probability of the new decay to that of the [decay](#) of B_c^+

into $J/\psi\pi^+$ has been calculated by various theorists over the last 30 years. Now these predictions can finally be compared with an experimental measurement: most predictions agree with the new result obtained by LHCb ($2.80\pm 0.15\pm 0.11\pm 0.16$).

The large number of b-quarks produced in LHC collisions and the excellent detector allows LHCb to study the production, decays and other properties of the B_c^+ meson in detail. Since the [meson](#)'s discovery by the CDF experiment at the Tevatron collider, 18 new B_c^+ decays have been observed (with more than five standard deviations), all of them by LHCb.

More information: Observation of the B_c^+ to $J/\psi\pi^+\pi^0$ decay, *arXiv* (2024). [DOI: 10.48550/arxiv.2402.05523](https://doi.org/10.48550/arxiv.2402.05523)

Provided by CERN

Citation: LHCb observes a new decay mode of the charmed beauty meson (2024, March 4) retrieved 27 April 2024 from <https://phys.org/news/2024-03-lhcb-decay-mode-charmed-beauty.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.