

First observation of photons-to-taus in proton–proton collisions

March 26 2024



Recreated candidate event of a $\gamma\gamma \rightarrow \tau\tau$ process in proton–proton collisions measured by the CMS detector. The tau can decay into a muon (red), charged pions (yellow) and neutrinos (not visible); energy deposits in the electromagnetic calorimeter in green and in the hadronic calorimeter in cyan. Credit: CMS collaboration.



In March 2024, the CMS collaboration announced the observation of two photons creating two tau leptons in proton–proton collisions. It is the first time that this process has been seen in proton–proton collisions, which was made possible by using the precise tracking capabilities of the CMS detector. It is also the most precise measurement of the tau's anomalous magnetic moment and offers a new way to constrain the existence of new physics.

The tau, sometimes called tauon, is a peculiar particle in the family of leptons. In general, leptons, together with quarks, make up the "matter" content of the Standard Model (SM). The tau was only discovered in the late 1970s at SLAC, and its associated neutrino—the <u>tau neutrino</u> —completed the tangible matter part upon its discovery in 2000 by the DONUT collaboration at Fermilab.

Precise research for the tau is rather tricky though, as its lifetime is very short: it remains stable for only $290 \cdot 10^{-15}$ s (a hundred-quadrillionth of a second).

The two other charged leptons, the electron and the muon, are rather well studied. A lot is also known about their magnetic moments and their associated anomalous magnetic moments. The former can be understood as the strength and orientation of an imaginary bar magnet inside a particle.

This measurable quantity, however, needs corrections at the quantum level arising from virtual particles tugging at the magnetic moment, deviating it from the predicted value. The quantum correction, referred to as <u>anomalous magnetic moment</u>, is of the order of 0.1%. If the theoretical and experimental results disagree, then this anomalous magnetic moment, a_l, opens doors to physics beyond the SM.

The anomalous magnetic moment of the electron is one of the most



precisely known quantities in particle physics and agrees perfectly with the SM. Its muonic counterpart, on the other hand, is one of the most investigated ones, into which research is ongoing. Although theory and experiments have mostly agreed so far, recent results give rise to a tension that requires further investigation.

For the tau, however, the race is still going. It is especially hard to measure its anomalous magnetic moment, a_{τ} , due to the tau's short lifetime. The first attempts to measure at after the tau's discovery came with an uncertainty that was 30 times higher than the size of the quantum corrections. Experimental efforts at CERN with the LEP and LHC detectors improved the constraints, reducing the uncertainties to 20 times the size of the quantum corrections.

In collisions, researchers look for a special process: two photons interacting to produce two tau leptons, also called a di-tau pair, which then decay into muons, electrons, or charged pions, and neutrinos. So far both ATLAS and CMS have observed this in ultra-peripheral lead–lead collisions. Now, CMS reports on the first observation of the same process during proton–proton collisions. These collisions offer a higher sensitivity to physics beyond the SM as new physics effects increase with the collision energy.

With the outstanding tracking capabilities of the CMS detector, the collaboration was able to isolate this specific process from others, by selecting events where the taus are produced without any other track within distances as small as 1 mm. "This remarkable achievement of detecting ultra-peripheral proton–proton collisions sets the stage for many groundbreaking measurements of this kind with the CMS experiment," said Michael Pitt, from the CMS analysis team.

This new method offers a new way to constrain the tau anomalous magnetic moment, which the CMS collaboration tried out immediately.



While the significance will be improved with future run data, their new measurement places the tightest constraints so far, with higher precision than ever before. It reduces the uncertainty from the predictions down to only three times the size of the quantum corrections.

"It is truly exciting that we can finally narrow down some of the basic properties of the elusive tau lepton," said Izaak Neutelings, from the CMS analysis team. "This analysis introduces a novel approach to probe tau g-2 and revitalizes measurements that have remained stagnant for more than two decades," added Xuelong Qin, another member of the analysis team.

A 3D interactive version of the event display with all tracks can be seen <u>here</u>.

Provided by CERN

Citation: First observation of photons-to-taus in proton–proton collisions (2024, March 26) retrieved 9 May 2024 from <u>https://phys.org/news/2024-03-photons-taus-protonproton-collisions.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.