

# Study shows how to increase success rate of restoration initiatives in the Cerrado

March 12 2024

---



Plots were sown with between zero and eight native grass species. The study was conducted in a former area of pasture taken over by the Chapada dos Veadeiros National Park (Goiás state, Brazil). Its purpose was to analyze the role of biodiversity in resistance to reinvasion of plant communities by exotic species.  
Credit: Guilherme Mazzochini

Statistics made available in January 2024 show that while deforestation in the Amazon fell by half in 2023, it increased by 43% in the Cerrado,

the Brazilian savanna. A research group at the State University of Campinas's Institute of Biology (IB-UNICAMP) in Brazil dedicated to ecological studies on the Cerrado has already [highlighted](#) the urgency of combating destruction of the biome. However, it is not enough to defend what remains: restoration is also needed.

Restoration presents a number of challenges, such as constant reinvasion by exotic grasses. Once these have been introduced into pasturelands, they spread vigorously, eliminate native species, and disfigure the herbaceous stratum, which is the Cerrado's biodiversity reserve and the mainstay of its ecosystem services.

They are hard to eradicate even after pasture is abandoned. The main plants in question are the tropical forage grasses *Urochloa* spp., *Melinis minutiflora* and *Andropogon gayanus*.

To validate ways of combating invasion or reinvasion by exotic grasses, a pioneering study was conducted by the IB-UNICAMP group at Chapada dos Veadeiros National Park in the state of Goiás, Center-West Brazil. The study by biologist Guilherme Mazzochini and collaborators [appears](#) in the *Journal of Applied Ecology*.

The study was part of the Thematic Project "Restoring Neotropical dry ecosystems—is plant functional composition the key to success?" for which the principal investigator is Rafael Oliveira, a professor at IB-UNICAMP. He is also the last author of the article. The penultimate author is Lucy Rowland, a professor at the University of Exeter in the UK.

"Many ongoing restoration projects aim to produce rapid plant cover in the area concerned, but in our study, we showed this isn't always the best

strategy. A large proportion of these projects lack a consistent scientific basis and are unsuccessful because they're unable to create biodiverse ecosystems that can withstand events such as drought, biological invasion, and fire."

"Any restoration initiative must be based on a profound understanding of the ecology of the ecosystem to be restored. Unfortunately, that's not usually the case," Oliveira said. He supervised Mazzochini's [postdoctoral research](#).

Biodiversity plays a key role in increasing resistance to invasion by exotic grasses, according to Oliveira. "Our findings show that higher diversity reduces invasion, probably owing to competition for resources in the nutrient-poor, acidic soil of the Cerrado," he said.

In conventional restoration projects, the typical management strategy deployed to eliminate exotic grasses before native species are planted involves controlled burning followed by frequent plowing during the dry season to remove stolons (above-ground creeping stems) and roots of exotic grasses, as well as to kill seedlings that have recently germinated.

However, the [exotic species](#) recolonize and dominate restored areas within a few years thanks to the large volume of seeds that remain below ground and the constant influx of seeds from the environs.

"Ecologists have observed for decades that species-rich local communities are more resistant to invasion than species-poor communities. Experiments with biodiversity and ecosystem functioning in regions with a temperate climate have confirmed these observations."

"In our experiment, plant communities with varying numbers of species were tested to understand how biodiversity influences productivity, nutrient cycling, and resistance to invasion, among other things. Our

study was the first to conduct an experiment involving biodiversity in the Cerrado and one of only a few ever conducted in the tropics," Mazzochini said.

Studies involving temperate ecosystems show that communities with more diversity tend to have lower levels of available resources, presumably owing to the complementary use of resources by species with different ecological strategies. This reduces invasion by exotic species, which generally require large amounts of resources to maintain rapid growth.

"Plants in any environment survive and grow by means of a range of ecological strategies associated with the anatomy and morphology of their leaves, stems, and roots. These characteristics are known as functional attributes," Mazzochini explained.

"Based on functional attributes, plant species can be classified on a spectrum from conservative to acquisitive. The most acquisitive species grow rapidly and have lighter, more tender vegetative organs that need more resources. These species, which include exotic grasses, are considered more competitive."

"On the other hand, conservative species grow more slowly and have a more robust structure with denser leaves, stems, and roots, so that they can withstand adverse conditions such as severe drought and low availability of nutrients."

Variation in functional attributes is used to classify species along the conservative axis, while ecologists recently introduced the label collaborative to identify adaptability, he explained. The concept centers on specific root length (SRL), defined as resource acquisition (absolute root length) divided by resource investment (mass).

"There's a direct correlation between SRL and root diameter. Thinner roots can absorb nutrients more autonomously and efficiently. Thicker roots often depend on symbiosis with mycorrhizal fungi, which establish mutualistic relationships with roots, gaining access to nutrients not available to plants, generally phosphorus, and in return, receiving the carbon produced by photosynthesis."

"This is a fascinating example of how plants interrelate with other organisms in order to prosper in many different environments," he said.

The two axes, conservative and collaborative, were considered in the study, which involved a biodiversity experiment conducted in an area of abandoned pasture taken over by Chapada dos Veados National Park.

"We installed 302 square plots measuring 2 m by 2 m and sowed zero to eight native grass species with a density of 1,000 seeds per square meter," he said.

The eight native species were *Axonopus aureus*, *Axonopus siccus*, *Andropogon fastigiatus*, *Aristida flaccida*, *Aristida riparia*, *Loudetiopsis chrysothrix*, *Schizachyrium sanguineum*, and *Trachypogon spicatus*.

As anticipated, a wave of reinvasion by exotic grasses occurred, reflecting the long period during which the area in question had been used as pasture. The invaders were *A. gayanus*, *M. minutiflora*, and three species of *Urochloa*. To assess the invasion of each plot, the researchers removed the exotic species and measured their biomass.

In line with the results of biodiversity experiments in temperate ecosystems, they found that communities with relatively high levels of species richness had less exotic biomass. Plots sown with only one native species had 3.6 times more exotic biomass than plots with eight species. This finding raised an important question: What was the mechanism responsible for conferring greater resistance to invasive species in more

species-rich communities?

"Our results suggest that more functional diversity in terms of plant height and SRL is associated with a reduction in invasion by exotic species. Height diversity, which relates to native species' above-ground biomass, appears to create several layers of vegetation, limiting the amount of sunlight that reaches invasive species."

"As we discovered, SRL diversity is a key factor, indicating that more varied root strategies and the shade provided by above-ground native biomass are effective in reducing invasion. Specifically, the monoculture plots had 4.7 times more exotic biomass on average than the plots with more SRL functional diversity," Mazzochini said.

This finding is especially important because many herbaceous communities in the Cerrado do not have large amounts of above-ground biomass, as the nutrient-poor, acidic soil limits productivity. Most of the biomass is stored below ground in roots and subterranean structures.

"We also compared the biomass of exotic species on monoculture plots and unseeded plots to see if the presence of native species alone facilitated invasion or competed with invasive species. This analysis showed that the effect of different native species varied."

"Some native species facilitated the establishment of exotic species, while others reduced invasion, even in the case of monoculture plots. We were surprised to find that the fastest-growing annual native species, *A. fastigiatus*, facilitated the establishment of exotic species, with 2.9 times more exotic biomass on monoculture than unseeded plots," Mazzochini said.

This finding could lead to a major revision of the restoration strategies adopted in the Cerrado, as *A. fastigiatus* is intensely used in restoration

projects. Given its rapid growth, the researchers expected it to cover the ground more quickly and prevent invasion by exotic species, but because it is an annual species, it dies early in the dry season, and its biomass decomposes and fertilizes the soil, favoring the establishment of fast-growing exotic species in subsequent years.

"To our surprise, the two species that most reduced invasion are the most different in functional terms. *S. sanguineum* is the smallest, with the longest SRL and the longest total root length. *A. siccus* is the tallest, with the shortest SRL and the shortest total root length. Plots where one of these species was the monoculture had 73% and 92% less biomass, respectively, than unseeded plots."

"These findings contradict the notion that a single native species can be sown with high density to prevent invasion. Instead, the results show the importance of using several ecological strategies when sowing [native species](#) for restoration purposes," Mazzochini said.

For Oliveira, biodiversity is key. "The greater the biodiversity, the greater the likelihood of defeating exotic species. The results also show the importance of revising management and restoration practices that use fast-growing annual or biennial species. Contrary to general belief, these species can facilitate the establishment of invaders over the years," he said.

**More information:** Guilherme G. Mazzochini et al, Effects of grass functional diversity on invasion success by exotic grasses in Cerrado grasslands, *Journal of Applied Ecology* (2023). [DOI: 10.1111/1365-2664.14561](https://doi.org/10.1111/1365-2664.14561)

Provided by FAPESP

Citation: Study shows how to increase success rate of restoration initiatives in the Cerrado (2024, March 12) retrieved 28 April 2024 from <https://phys.org/news/2024-03-success-cerrado.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.