

Advance in forensic fingerprint research provides new hope for cold cases

April 22 2024, by Meg Cox



Test sample preparation of (a) ZPD doped fingerprint and (b) fresh fingerprint: (1) wash and dry hands, put on nitrile gloves, wear for 1 h, turn inside-out and put back on; (2) gently touch ZPD powder with a gloved finger; (3) generate a fingermark onto a glass slide containing (a) ZPD residue or (b) no ZPD residue; (4) take-up the fingerprint from the slide surface using a piece of gel-lifter; and (5) store the gel-lifter piece in a dark place at room temperature ($22 \pm 1^{\circ}$ C) (created with Biorender.com). Credit: *Drug Testing and Analysis* (2024). DOI: 10.1002/dta.3688



Researchers have unveiled a method capable of detecting drug substances from fingerprints lifted from crime scenes, which could provide fresh insights into unsolved cases. The research is <u>published</u> in the journal *Drug Testing and Analysis*.

Analytical scientists from Loughborough University have demonstrated for the first time that <u>drug</u> residue—namely the fast-acting sleeping pill Zolpidem, which has been <u>linked to drug-facilitated sexual assault and</u> <u>drink spiking</u>—can be detected on gel-lifted fingerprints.

Dr. Jim Reynolds and Dr. Ayoung Kim say the breakthrough could shed new light on cold cases and unsolved crimes as forensic gel lifters—which transfer prints onto a gelatin surface—are used globally by scenes of crimes officers to preserve and visualize fingerprints.

"This is the first time that analysis of gel-lifted prints for a drug substance has been accomplished, and shows that lifted prints and other forensic marks can be interrogated for useful information," says Dr. Reynolds, the research lead.

"Since gel-lifted prints and marks can be stored for many years, the technique could be of real use in cold cases where additional information may prove useful to either link or exonerate a suspect to the investigation. Working with <u>police forces</u> and applying the method to cold case samples could help bring criminals to justice who may have thought they have got away with it."

A number of tests exist to detect drugs directly from fingerprints, but these face limitations. They can be destructive to the fingerprint, degrade drug residues, and be affected by environmental interferences.



It has long been speculated that gel-lifted prints contain valuable chemical information and could offer more accurate drug detection.

However, traditional techniques used to analyze the chemicals present in a sample have previously not been suitable for gel lifters. This is because they detect all chemicals present, including those that make up the gel, making it difficult to identify specific substances.

The method used by Dr. Reynolds and Dr. Kim, called sfPESI-MS, overcomes this issue using a rapid separation mechanism that distinguishes the drug substance from the background of the gel.

The process involves sampling the chemicals from the gel lifters into tiny liquid droplets. The chemicals extracted into the droplets are then ionized, which means they gain or lose electric charge depending on their chemical properties. The drug substance chemicals are more surface active than the chemicals originating from the gel, which enables them to be separated from the mixture.

This separation method enables the direct detection of a drug substance using <u>mass spectrometry</u>, a technique that identifies chemicals by measuring their molecular weight. The researchers have successfully tested the technique using Zolpidem-laced fingerprints lifted from glass, metal, and paper surfaces in a laboratory setting.

They now hope to work with police forces to analyze stored gel-lifted prints and use the method to identify other substances.

Dr. Reynolds said, "Zolpidem was the focus of our research, but the method could just as easily be applied to other drug substances a person may have been handling and could be applied to other chemicals such as explosives, gunshot residues, paints, and dyes.



"By linking chemical information to the fingerprint, we can identify the individual and link to the handling of an illicit substance which may prove useful in a prosecution. This could be useful to detect individuals who have been spiking drinks; for example, if the drug they are using gets onto their fingertips, then they will leave evidence at the scene."

Dr. Kim, who is the first author of the paper and completed the research as part of her Ph.D. at Loughborough, added, "We would like to apply our method to real samples from criminal investigations; it would be good to know my Ph.D. research has helped bring criminals to justice."

More information: Ayoung Kim et al, A direct analysis method using sheath flow probe electrospray ionisation-mass spectrometry (sfPESI-MS) to detect drug residues from fingerprint forensic gel lifts, *Drug Testing and Analysis* (2024). DOI: 10.1002/dta.3688

Provided by Loughborough University

Citation: Advance in forensic fingerprint research provides new hope for cold cases (2024, April 22) retrieved 17 May 2024 from https://phys.org/news/2024-04-advance-forensic-fingerprint-cold-cases.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.