

Airborne interferometric radar altimeter shows potential for submesoscale sea surface height anomaly measurements

April 23 2024, by Zhang Nannan



Comparison of the SSHA wavenumber spectrum observed by AIRA (in red)



with the SSHA wavenumber spectra from satellite altimeters ICESat-2, SARAL/AltiKa, and Sentinel-6A. Credit: Xu Yongsheng

A research team led by Prof. Xu Yongsheng from the Institute of Oceanology of the Chinese Academy of Sciences has obtained the twodimensional sea surface height anomaly (SSHA) of about 100 km along the track and the first broadband SSHA wavenumber spectrum using Airborne Interferometric Radar Altimeter (AIRA) observations.

Their study was **<u>published</u>** in *Remote Sensing* on April 12.

To date, humanity has not been able to observe two-dimensional (2D) oceanic processes at the 0.1–10 km submesoscale in the spatial domain using remote sensing. The SSHA signal at this scale is small and exceeds the resolution limits of the satellite altimeters used to date.

However, oceanic processes at this scale play a critical role in the study of ocean energy transfer, cascading, and dissipation, and are crucial for research on ocean energy balance, nutrient transport, and global climate change studies.

In this study, the researchers provided a detailed analysis of the SSHA and its wavenumber spectrum obtained by AIRA, and demonstrated the potential of AIRA for submesoscale SSHA observations. The AIRA shows a good ability to capture small-scale SSHA. In the boxed section of the image below, waves with wavelengths between 50 m and 150 m stand out as the most dominant feature. The submesoscale SSHA typically has much smaller magnitudes than <u>ocean waves</u>.





2D SSHA from the IRA measurements. Credit: Xu Yongsheng

Nevertheless, it is clear that the waves are superimposed on a background characterized by undulating heights. AIRA's spectrum has a distinct negative slope down to about 1 km, consistent with theoretical predictions.

AIRA's resolved power spectral density level in the 1–10 km submesoscale range is about ten times lower than that of ICESat-2 ATL03, demonstrating AIRA's superior ability to resolve lower energy SSHA variations. The wavenumber spectrum is an important basis for evaluating the observational performance of AIRA.

"The results of this study highlight the tremendous potential of AIRA in detecting SSHA at a range of scales from oceanic waves to submesoscale," said He Jinchao, first author of the study.

"We expect that our study will inspire a new appreciation for the potential of AIRA in detecting two-dimensional small submesoscale SSHA," added Prof. Xu, corresponding author of the study. "These processes are critical to understanding ocean dynamics, <u>marine</u>



ecosystems, and their role in the Earth's climate system."

More information: Jinchao He et al, Sea Surface Height Wavenumber Spectrum from Airborne Interferometric Radar Altimeter, *Remote Sensing* (2024). DOI: 10.3390/rs16081359

Provided by Chinese Academy of Sciences

Citation: Airborne interferometric radar altimeter shows potential for submesoscale sea surface height anomaly measurements (2024, April 23) retrieved 17 May 2024 from <u>https://phys.org/news/2024-04-airborne-interferometric-radar-altimeter-potential.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.