



Ant pheromones may help protect hikers and campers from ticks

April 11 2024

(a)



(b)



Drawings illustrating (a) the location of the poison gland and Dufour's gland in *Formica oreas* worker ants, and (b) the olfactometer used in tick bioassays. For bioassays, the lateral chambers of the olfactometer received a piece of filter paper treated with a treatment or control stimulus, and a damp cotton ball to increase relative humidity. A single bioassay tick was released into the central chamber, and was considered a responder, if it was found at the end of the bioassay in a lateral chamber, or in a connecting glass tube closer to a lateral chamber than to the central chamber. PG, poison gland; PGR, poison gland reservoir; DFG, Dufour's gland; CR, crop; MG, midgut; HG, hindgut. Credit: *Royal Society Open Science* (2024). DOI: 10.1098/rsos.231355

If you're outside enjoying the spring sunshine, then chances are, ticks are too. However, new research from Simon Fraser University (SFU) suggests the use of ant pheromones as a topical repellent, or as an environmental barrier, may help protect hikers and campers from the harmful bites of black-legged ticks, which could carry Lyme disease.

"Ticks like a lot of the places and a lot of the times of years that we also like to be outside enjoying the weather," says lead author Claire Gooding, a member of the SFU Gries Lab. "There's quite a big risk of tick encounters in the summer, outdoor sports season. People often encounter them on the sides of trails."

A paper [published](#) in *Royal Society Open Science* looked at what predators (ants, spiders or beetles) black-legged ticks might avoid and what adaptations they might have to avoid them.

"We decided to look at ants because they are [social insects](#) and use a huge range of pheromones to communicate with one another," says Gooding. "They're chemically noisy. And for something that perceives the world chemically, they're easy to predict where they'll be, based on these pheromones."

Ticks transmit more diseases than any other arthropods and their natural ranges may be expanding due to [climate change](#), increasing the need for management strategies as well as new repellents. The study found that ticks avoided surfaces where ants had been, even if those ants were removed.

"They could see that there were ants and basically go, 'I'm not going to go there, because there may be ants there, or there may be ants there again soon in the future.'"

From there, Gooding and the team identified the specific chemical pheromones as well as the two ant glands that produced them. They then worked with a synthetic chemist and an analytical chemist to recreate a synthetic version of those pheromones. They found that ticks avoided the synthetic pheromone too.

The team, which works with industrial sponsors to further discoveries into the applied world, has a [patent application](#) on the repellent chemicals. They are hoping to bring them to market for real-world use, either as a topical repellent (like [mosquito repellent](#)), or as an environmental repellent, which could be used with a physical barrier, like [wood chips](#), to deter [ticks](#) from entering an area or a hiking trail.

More information: Claire E. Gooding et al, Blacklegged ticks, *Ixodes scapularis*, reduce predation risk by eavesdropping on communication signals of Formica ants, *Royal Society Open Science* (2024). [DOI: 10.1098/rsos.231355](https://doi.org/10.1098/rsos.231355)

Provided by Simon Fraser University

Citation: Ant pheromones may help protect hikers and campers from ticks (2024, April 11)
retrieved 20 May 2024 from <https://phys.org/news/2024-04-ant-pheromones-hikers-campers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.