

Astronomers find 10 new millisecond pulsars in globular cluster Terzan 5





Integrated pulse profiles of all the newly discovered pulsars in Terzan 5. These plots are obtained after summing together individual profiles from different epochs and aligning the profiles with respect to a reference template profile. Credit: Padmanabh et al., 2024.

Using the MeerKAT radio telescope, an international team of



astronomers has detected 10 new millisecond pulsars in a Galactic globular cluster known as Terzan 5. The finding, which makes Terzan 5 the most pulsar-rich globular cluster, was reported in a research <u>paper</u> published March 26 on the pre-print server *arXiv*.

Pulsars are highly magnetized, rotating <u>neutron stars</u> emitting a beam of electromagnetic radiation. The most rapidly rotating pulsars, with rotation periods below 30 milliseconds, are known as <u>millisecond pulsars</u> (MSPs). Astronomers assume that they are formed in binary systems when the initially more massive component turns into a neutron star that is then spun up due to accretion of matter from the secondary star.

Discovered in 1968, Terzan 5 is a globular cluster (GC) located in the Milky Way's bulge, some 18,800 <u>light years</u> away. It has a radius of 2.7 light years, mass of about 2 million solar masses, and is estimated to be 12 billion years old. Given that Terzan 5 has been known for decades, it is one of the most widely studied GCs, and previous observations have found that it hosts a numerous population of pulsars, including MSPs.

The presence of many pulsars in Terzan 5 may still be undisclosed and it is estimated that this cluster may contain even 200 such objects. Therefore, a team of astronomers led by Prajwal Padmanabh of the Max Planck Institute for Gravitational Physics in Hannover, Germany, has observed Terzan 5 with the MeerKAT radio telescope using the L-Band (856–1,712 MHz) receiver, hoping to identify new pulsars in this cluster.

As a result, Padmanabh's team detected 10 new MSPs, out of which nine turned out to be binary systems. After the discovery, Terzan 5 now hosts 49 pulsars, which is so far the highest number for any known globular cluster.



The newfound MSPs have spin periods between 1.25 and 4.54 milliseconds, while their dispersion measures are similar—within a range of 235.44–239.46 pc/cm³. The astronomers managed to derive the characteristic age for five of the newly detected pulsars and it is generally between 1.3 and 3.18 billion years.

The researchers underlined that the only isolated pulsar in the sample, designated Ter5as, has a spin period of 2.32 milliseconds and many of its parameters still remain unknown. Four pulsars in the sample, namely Ter5au, Ter5ap, Ter5av, and Ter5ax likely have white dwarf companions with minimum masses within a range of 0.07–0.28 solar masses.

The authors of the paper noted that one of the newly found MSPs, designated Ter5ao, is an eccentric, wide-orbit <u>pulsar</u> with a large minimum companion mass of approximately 0.69 <u>solar masses</u> and a rapid spin period. These findings suggest that it may be a rare double neutron star (DNS) system.

More information: P. V. Padmanabh et al, Discovery and timing of ten new millisecond pulsars in the globular cluster Terzan 5, *arXiv* (2024). DOI: 10.48550/arxiv.2403.17799

© 2024 Science X Network

Citation: Astronomers find 10 new millisecond pulsars in globular cluster Terzan 5 (2024, April 2) retrieved 20 May 2024 from <u>https://phys.org/news/2024-04-astronomers-millisecond-pulsars-globular-cluster.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.