


# Boron deficiency in oilseed rape transcriptome resembles a wounding and infection response

April 15 2024, by Magdalena Eisenmann



Overview of phenotypic effects of B-deficient (-B) and B-sufficient (+B) growth conditions on the overall and inflorescence development of *Brassica napus* at (A) growth stage BBCH 51 (main inflorescence in the middle of the top leaves visible from above, flowers not open), (B) growth stage BBCH 60–61 (first flowers open), (C) growth stage BBCH 60–61 plus 4 days, and (D) growth stage BBCH 60–61 plus 14 days. Credit: *Physiologia Plantarum* (2023). DOI: 10.1111/ppl.14088

Boron deficiency has a devastating effect on oilseed rape and related plants. However, little is known about the underlying genetic mechanisms. A study shows that the response to persistent or short-term acute boron deficiency is similar to that seen in the response to pests and infections. These results lay the foundation for breeding plants that can better cope with boron deficiency and for avoiding related yield losses.

Oilseed rape and related plants have a high boron requirement. A boron deficiency is often invisible on the outside of the affected plants for a long time, especially during vegetative stages. Nevertheless, it has serious consequences: First, the deficiency can inhibit root growth, and then later, the flowers can shrivel and die prematurely. This can result in enormous yield losses, especially when seeds or grains are the crop product of interest.

In order to understand the underlying mechanisms, researchers from the Technical University of Munich (TUM), Bielefeld University and the Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) have investigated which genes are active when boron is deficient and which are active when there is sufficient supply.

Their study is [published](#) in the journal *Physiologia Plantarum*.

In a first step, the plants react to boron deficiency by producing more transport proteins that take up the nutrient. If deficiency persists, though, a stress reaction comes to the foreground in the developing and yield-relevant flowers. This resembles the reaction to pest attacks, wounding or infections and leads to the death of the affected flowers.

It is still not known whether the response is a targeted reaction of the plant to save resources for a potential later flowering or whether the [signaling pathway](#) is triggered unintentionally. Both boron deficiency and infections damage plant cell walls. Cell wall damage could mimic

infection and lead to a similar signal even in the absence of an infectious threat.

## **Boron-efficient plant varieties secure yields**

Climate change could make boron shortages even more frequent in the future. In Central Europe, winters with more precipitation and prolonged dry periods in spring and summer are expected. As a result, water-soluble nutrients will be increasingly washed out in winter. In spring, the soil lacks moisture for the plants to absorb the nutrients.

Under these conditions, it is difficult to compensate for the nutrient deficiency through targeted fertilization. "Either the boron in the soil does not reach the plant, or an already undetected boron deficiency leads to damage to the pathways for water and nutrients. As a result, applied boron can no longer be transported to the places in the plant where it is needed," says Prof. Gerd Patrick Bienert from the TUM Chair of Crop Physiology.

Developing plant varieties that can either handle boron efficiently or are both water and boron efficient can secure future yields. The researchers' findings contribute to identifying possible starting points in plant breeding.

"The identification and breeding of boron efficiency mechanisms and the cultivation of boron-efficient varieties will become increasingly important in the context of climate change. A sufficient supply of [boron](#) is essential for the yield stability of crops, especially when water is scarce," says crop researcher Bienert.

**More information:** Bart Verwaaijen et al, The Brassica napus boron

deficient inflorescence transcriptome resembles a wounding and infection response, *Physiologia Plantarum* (2023). DOI: [10.1111/ppl.14088](https://doi.org/10.1111/ppl.14088)

Provided by Technical University Munich

Citation: Boron deficiency in oilseed rape transcriptome resembles a wounding and infection response (2024, April 15) retrieved 16 June 2024 from <https://phys.org/news/2024-04-boron-deficiency-oilseed-rape-transcriptome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.