

Researchers clarify cryptic differentiation within enigmatic hemiparasitic love vine *Cassytha filiformis*

April 15 2024, by Zhang Nannan



Cassytha filiformis (flowers). Credit: [Forest & Kim Starr](#)/Wikimedia Commons, [CC BY](#)

The widespread hemiparasitic Lauraceae genus *Cassytha* currently contains 19 described species, one variety and four forms. This genus is controversial and has not been satisfactorily resolved. *Cassytha filiformis* is cosmopolitan in tropical and subtropical regions. It is currently the only species reported from China.

Accurate [species](#) delimitation has been a great challenge for biodiversity conservation practice, and many species may have already gone extinct before being recognized and described, especially for morpho-hyperspecialized parasitic, hemiparasitic plants, where it is very difficult to accurately capture the species.

In a study [published](#) in the *Journal of Systematics and Evolution*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) of the Chinese Academy of Sciences and their collaborators examined Chinese *Cassytha* samples that are currently placed in *C. filiformis*, as well as several related members of the racemose group, to investigate the phylogenetic and evolutionary history of the genus in the region.

Using integrated means such as phylogenetic analysis, morphological trait statistical analysis, and species distribution simulation, the researchers investigated whether there are cryptic species of *Cassytha filiformis* within China and neighboring regions.

The results of the [phylogenetic analysis](#) showed that *C. filiformis* formed two well-supported monophyletic lineages, and the morphological analyses revealed that the two lineages differed significantly in both inflorescence stem thickness and fruit type ratio. Species distribution simulations showed that different environmental factors influenced their distributions.

Moreover, the divergence time of the two lineages was inferred to be around the Early Miocene, suggesting that the divergence of the cryptic

hemiparasitic love vine species may have been related to new ecological opportunities and host diversification brought about by the East Asian monsoon in the Early Miocene.

"Our study revealed the cryptic diversity of the enigmatic hemiparasitic love vine *Cassytha filiformis* in China and neighboring areas. It provides new evidence for the widespread existence of cryptic species in parasitic plants," said Li Jie of XTBG.

More information: Zhi-Fang Liu et al, Dense infraspecific sampling reveals cryptic differentiation in the enigmatic hemiparasitic love vine *Cassytha filiformis* (Lauraceae), *Journal of Systematics and Evolution* (2024). [DOI: 10.1111/jse.13069](https://doi.org/10.1111/jse.13069)

Provided by Chinese Academy of Sciences

Citation: Researchers clarify cryptic differentiation within enigmatic hemiparasitic love vine *Cassytha filiformis* (2024, April 15) retrieved 21 May 2024 from <https://phys.org/news/2024-04-cryptic-differentiation-enigmatic-hemiparasitic-vine.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|