

Deep parts of Great Barrier Reef 'insulated' from global warming, for now

April 8 2024



Photos taken before and after bleaching in the northern Great Barrier Reef near Lizard Island. Credit: Dr. George Roff

Some deeper areas of the Great Barrier Reef are insulated from harmful heat waves—but that protection will be lost if global warming continues, according to new research.

High surface temperatures have caused mass "bleaching" of the Great Barrier Reef in five of the last eight years, with the latest happening now.

Climate change projections for [coral reefs](#) are usually based on [sea](#)

[surface temperatures](#), but this overlooks the fact that deeper water does not necessarily experience the same warming as that at the surface.

A new study—led by the universities of Exeter and Queensland—examined how changing temperatures will affect mesophotic corals (depth 30–50 meters). The paper, published in the journal *Proceedings of the National Academy of Sciences*, is titled, "Climate change impacts on mesophotic regions of the Great Barrier Reef."

It found that separation between warm buoyant surface water and cooler deeper water can insulate reefs from surface heat waves, but this protection will be lost if [global warming](#) exceeds 3°C above pre-industrial levels.

The researchers say similar patterns could occur on other reefs worldwide, but local conditions affecting how the water moves and mixes will mean the degree to which deeper water coral refuges exist and remain insulated from surface heat waves will vary.

"Coral reefs are the canary in the coalmine, warning us of the many species and ecosystems affected by climate change," said Dr. Jennifer McWhorter, who led the research during a QUEX Ph.D. studentship at the universities of Exeter and Queensland.

"Coral bleaching is a dramatic sign of the impact humans are having on the planet.

"Our study offers both hope and a warning—hope that some reefs are resilient to current levels of climate change, and a warning that this resilience has its limits."



Photos taken before and after bleaching in the northern Great Barrier Reef near Lizard Island. Credit: Dr. George Roff

The study finds that 3°C of global warming would push mesophotic temperatures in the Great Barrier Reef past 30°C —a recognized threshold for coral mortality.

This does not necessarily mean that all coral would die, but it would place the reef in a state of stress that would increase mortality and possibly cause it to collapse.

Dr. McWhorter, now at NOAA's Atlantic Oceanographic & Meteorological Laboratory, said, "Some shallow-water species are not found in deeper areas—so mesophotic reefs can't provide refuges for them as shallow reefs are degraded.

"And, as our study shows, mesophotic corals are themselves threatened if global warming continues."

To calculate their projections of mesophotic reefs warming, the research

team considered factors such as wind and tidal mixing of water, and local complexities.

They estimate that, by 2050–60, bottom temperatures on the Great Barrier Reef (30–50 meters) will increase by 0.5–1°C under lower projected greenhouse gas emissions, and 1.2–1.7°C under higher emissions.

Dr. Paul Halloran, from Exeter's Global Systems Institute, said, "To protect coral reefs, we need to understand them better.

"Reefs face multiple threats—not just climate change. By targeting management of these threats on reefs that have the best chance of escaping the worst impacts of climate change, hopefully some healthy reefs can be maintained.

Professor Peter Mumby, from the University of Queensland, said, "There is so much to learn about deeper, tropical coral reefs, especially as we cannot assume that their depth provides a persistent refuge from the consequences of rising global carbon emissions."

More information: McWhorter, Jennifer K., Climate change impacts on mesophotic regions of the Great Barrier Reef, *Proceedings of the National Academy of Sciences* (2024). [DOI: 10.1073/pnas.2303336121](https://doi.org/10.1073/pnas.2303336121). doi.org/10.1073/pnas.2303336121

Provided by University of Exeter

Citation: Deep parts of Great Barrier Reef 'insulated' from global warming, for now (2024, April 8) retrieved 2 May 2024 from <https://phys.org/news/2024-04-deep-great-barrier-reef-insulated.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.