

Diverse native wildflower plantings for pollinators in farmlands

April 16 2024



Pollinators are declining rapidly, largely due to land conversion and

intensification of agriculture. To mitigate their crisis, low-disturbance habitats, such as sown wildflower plantings, could promote pollinators by restoration of their resources. However, comprehensive knowledge is lacking on how landscape context, spatial configuration and age of wildflower plantings, seasonality and flower composition affect pollinator communities, especially from East-Central Europe. To understand these effects, researchers of HUN-REN Centre for Ecological Research established diverse native wildflower plantings within heterogeneous and homogeneous agricultural landscapes. Credit: Borbála Bihaly (top left, bottom right) and Áron Bihaly (bottom left, middle and top right)

Pollinators are declining rapidly, largely due to land conversion and intensification of agriculture. To mitigate their crisis, low-disturbance habitats, such as sown wildflower plantings (commonly known forms are wildflower strips at the edges of arable fields), could promote pollinators by restoration of their resources (food, sheltering and nesting habitats). However, comprehensive knowledge is lacking on how landscape context, spatial configuration and age of wildflower plantings, seasonality and flower composition affect pollinator communities, especially from East-Central Europe.

To understand these effects, researchers from the HUN-REN Centre for Ecological Research have established diverse native wildflower plantings within heterogeneous and homogeneous agricultural landscapes, in two spatial configurations: one large field or three smaller strips. Floral resources and wild pollinator insects ([wild bees](#), hoverflies, butterflies) were sampled, in early and mid-summer, for two years after establishment (2020-21).

The team [reports](#) on their work in *Agriculture, Ecosystems & Environment*.

Flower resources of the sown [plant species](#) increased continuously, and

were complemented at high rate by flowering plant species from the soil seed bank, especially in the first year. Both flower abundance and diversity increased the abundance of pollinators, highlighting the important role of using diverse seed mixtures. Wild bee abundance and [species richness](#) increased year by year and season by season, while butterfly abundance also demonstrated a yearly increase after establishment.

Hoverfly abundance and species richness, however, showed an opposite trend, possibly due to the inter-annual variation. Wild bee and butterfly abundance was higher in the heterogeneous than in the homogeneous landscapes. Researchers did not observe any significant local effects of spatial configuration itself on pollinator populations.

Wildflower plantings were particularly attractive for wild bees in flower-poor (homogeneous) landscapes during the flower-poor (mid-summer) period of the year. This can be explained by the increased contrast in flower resources between the wildflower plantings and the surrounding landscape, and highlights the importance of wildflower plantings in such conditions.



Researchers of HUN-REN Centre for Ecological Research established diverse native wildflower plantings within heterogeneous and homogeneous agricultural landscapes to mitigate pollinator's crisis. Credit: Viktor Szigeti (top left and middle left) and Borbála Bihaly (bottom row, top right and middle right)

The results emphasize that to support pollinators effectively, future wildflower plantings should be maintained for multiple years, in order to maximize floral diversity and ensure continuously available flower resources throughout the entire season. It can be facilitated by using diverse seed mixtures of native flowering plants; particularly focusing on plants that bloom during the crucial mid-summer period; allowing the plant species to grow and bloom from the soil seed bank, and using management techniques, which help to extend the flowering season, said Áron Bihaly, corresponding author of the publication.

Further results from the upcoming years and similar long-term and landscape-scale experimental studies are needed to understand all the benefits and ecological processes of diverse native wildflower plantings especially in understudied European regions.

More information: Á.D. Bihaly et al, Landscape-wide floral resource deficit enhances the importance of diverse wildflower plantings for pollinators in farmlands, *Agriculture, Ecosystems & Environment* (2024). [DOI: 10.1016/j.agee.2024.108984](https://doi.org/10.1016/j.agee.2024.108984)

Provided by Ökológiai Kutatóközpont

Citation: Diverse native wildflower plantings for pollinators in farmlands (2024, April 16) retrieved 20 May 2024 from <https://phys.org/news/2024-04-diverse-native-wildflower-pollinators-farmlands.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
