

# East coast mussel shells are becoming more porous in warming waters

April 17 2024

---



A selection of historic and modern mussel shells from the American Museum of Natural History's collection used in the study. Credit: Daniel Kim/ AMNH

Researchers at the American Museum of Natural History have found that over the last 120 years, the porosity—or small-scale holes—in mussel shells along the East Coast of the United States has increased, potentially due to warming waters. The study, which analyzed modern mussel shells in comparison to specimens in the Museum's historic collection, was [published](#) in the journal *PLoS ONE*.

"Mussels are important on so many levels: They provide habitats on reefs, they filter water, they protect coasts during storms, and they are important commercially as well—I love mussels and I know many other people do, too," said Leanne Melbourne, a Kathryn W. Davis postdoctoral fellow in the Museum's Master of Arts in Teaching program and the lead author on the study. "Human-caused [environmental changes](#) are threatening the ability of mussels and other mollusks to form their shells, and we need to better understand what risks will come from this in the future."

Previous studies on the blue (common) mussel (*Mytilus edulis*) have used [lab experiments](#) to investigate how ocean warming and acidification impact the form and function of these important marine invertebrates. Melbourne drew on a historical resource—the Museum's large bivalve collection, which dates to the late 1800s—to trace environmental changes in natural settings.

"Museum collections are ideal for showing change over long stretches of time. Organisms in natural environments have slow exposure to changing conditions, with more time to adapt and more than one stressor," said co-author Nathalie Goodkin, a curator in the Museum's Department of Earth and Planetary Sciences. "Laboratory experiments are great at teasing apart individual drivers, but that's not how things work in the real world."

Melbourne focused on shells collected in the early 1900s and the 1960s from five sites: Nahant Bay in Massachusetts, the southern end of Cape Cod, the tip of the Long Island Sound, western Long Island Sound, and New York Harbor. She then paired these historic specimens with modern samples from the same sites and conducted a series of comparisons including the thickness of the shells, their surface area and volume, and density and porosity.



The blue (common) mussel (*Mytilus edulis*). Credit: Daniel Kim/ AMNH



Lead author Leanne Melbourne collecting mussel shells on Canoe Beach in Nahant Bay, Massachusetts. Credit: Maria Rosabelle Ong



Lead author Leanne Melbourne with a selection of the modern and historic mussel shells used in the new study. Credit: Daniel Kim/ AMNH

The study found that shells collected today are significantly more porous than shells collected in the 1960s and more porous than shells collected at some of the sites in the early 1900s, suggesting that increases in temperature led to increases in porosity.

The North Atlantic has experienced seasonal temperature increases of up to 3 degrees Celsius (5.4 degrees Fahrenheit) since 1902, but the

warming has not been uniform, which may account for some of the variations in porosity in historical specimens. The influence of the North Atlantic Oscillation, a weather phenomenon that brought colder-than-expected water temperatures in the 1960s, is also a factor.

Porosity is known to affect the structural integrity of mollusk shells, making shells weaker and potentially more susceptible to damage. More work is needed to determine how [temperature change](#), in particular, could be affecting these mussel populations.

"We know that mussel and [mussel](#) reef ecosystem services are dependent on robust [shell](#) formation. If they are forming weaker shells, they will break more easily, and we might lose these important organisms," Melbourne said. "But we need more studies first to look at what's happening here on a materials science level."

**More information:** Leanne A. Melbourne et al, Using Museum collections to assess the impact of industrialization on mussel (*Mytilus edulis*) calcification, *PLoS ONE* (2024). [DOI: 10.1371/journal.pone.0301874](#)

Provided by American Museum of Natural History

Citation: East coast mussel shells are becoming more porous in warming waters (2024, April 17) retrieved 2 May 2024 from <https://phys.org/news/2024-04-east-coast-mussel-shells-porous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.