


# Electro-optic 3D snapshot of a laser wakefield accelerated kilo-ampere electron bunch

April 9 2024



Relativistic electron bunches were generated via the incidence of a high power laser onto a gas jet. The TR radiation emitted when electron passing through a metal foil was imaged to an EO crystal. OTR imaging and EO spatial decoding were conducted simultaneously to achieve the electron spatial-temporal profile in a single shot. Credit: Kai Huang, Zhan Jin, Nobuhiko Nakanii, Tomonao Hosokai, and Masaki Kando

Invented by T. Tajima and J. Dawson, laser wakefield acceleration (LWFA) harnesses the power of high-intensity laser pulses to drive plasma waves with acceleration gradients orders of magnitude higher than conventional RF-based accelerators.

Treating the [laser](#) pulse as a bullet, the ponderomotive force induces a periodic wave in plasma, akin to an RF cavity. Electrons trapped in this wave can reach energies of GeV within a mere centimeter—a thousand times shorter than conventional accelerators. The femtosecond temporal durations of electron bunches from LWFA present unprecedented opportunities for ultrafast studies.

Over the past two decades, LWFA has seen remarkable advancements in maximum energy, energy spread, charge, and repetition rate.


With its high acceleration gradient and ability to produce high-brightness electron bunches, LWFA holds immense potential for applications in high-energy physics, X-ray pump-probe studies, and time-resolved dosimetry. The three-dimensional (3D) density is a critical parameter influencing the luminosity in a collider, the brightness of secondary X-ray sources, and the peak dose rates of radiation, respectively.

In particular, a tabletop X-ray free-electron laser (XFEL) is intricately linked to the lasing process in an undulator. However, until now, experimental measurement has remained elusive due to the lack of diagnostic methods. While transverse or relative longitudinal distributions have been studied separately, the absolute 3D density profile has yet to be fully characterized.

In a new paper [published](#) in *Light: Science & Applications*, a team of scientists, led by Dr. Masaki Kando from Kansai Institute for Photon Science (KPSI), National Institutes for Quantum Science and Technology (QST), Japan, Prof. Tomonao Hosokai from SANKEN,

Osaka University, Japan, and co-workers have achieved a breakthrough in the single-shot monitoring of the 3D density profile of LWFA electron bunches.

Through experimental and numerical investigations on the optical transition radiation (OTR) imaging, electro-optic (EO) spatial decoding, and genetic algorithm (GA), the detailed 3D structures of the ultrafast electron bunches from LWFA were successfully reconstructed, illuminating the intricate beam dynamics of the electron bunch.



With the OTR and EO signals, the electron 3D density profile was reconstructed by using the genetic algorithm. Credit: Kai Huang, Zhan Jin, Nobuhiko Nakanii, Tomonao Hosokai, and Masaki Kando

The findings reveal a transverse size of less than 30 micrometers for the electron bunch, showing the remarkable resolution achieved by the OTR imaging technique. Moreover, the current profile exhibited a complex multi-peak shape, with a sub-10 femtosecond structure boasting a peak current exceeding 1 kiloampere (kA)—a testament to the exceptional

performance of laser wakefield acceleration.

Of particular significance is the observed peak 3D number density of  $\sim 9 \times 10^{21} \text{ m}^{-3}$ , offering invaluable insights into the accelerated electron bunch. This detection showed the potential for implementing a detector at any position along a beam transport line, thus opening new avenues for future applications in accelerator science and beyond.

"The ability to measure the three-dimensional density profile of electron bunches will enhance our understanding of LWFA and unlock its full potential for diverse applications," said Dr. Kai Huang, senior researcher at QST and the lead author of the paper.

"The results and methodologies presented in this article have far-reaching implications across a spectrum of disciplines, including accelerator physics, high-power lasers, and terahertz optics."

**More information:** Kai Huang et al, Electro-optic 3D snapshot of a laser wakefield accelerated kilo-ampere electron bunch, *Light: Science & Applications* (2024). [DOI: 10.1038/s41377-024-01440-2](https://doi.org/10.1038/s41377-024-01440-2)

Provided by Changchun Institute of Optics, Fine Mechanics And Physics

Citation: Electro-optic 3D snapshot of a laser wakefield accelerated kilo-ampere electron bunch (2024, April 9) retrieved 2 May 2024 from <https://phys.org/news/2024-04-electro-optic-3d-snapshot-laser.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.