


Evidence of a new subatomic particle observed

April 12 2024



A portrait of nucleon-antinucleon bound state. Credit: *Physical Review Letters* (2024). DOI: 10.1103/PhysRevLett.132.151901


The BESIII collaboration have reported the observation of an anomalous line shape around $p\bar{p}$ mass threshold in the $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ decay, which indicates the existence of a $p\bar{p}$ bound state. [The paper](#) was

published online in *Physical Review Letters*.

The proximity in mass to $2m_p$ is suggestive of nucleon-antinucleon bound states, an idea that has a long history. Before the birth of Quark Model, a nucleon-antinucleon bound state was already proposed by Prof. E. Fermi and Prof. C. N. Yang.

There is an accumulation of evidence for anomalous behavior in the proton-antiproton system near the $p\bar{p}$ mass [threshold](#), e.g., $J/\psi \rightarrow \gamma p\bar{p}$, $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$ and the proton's effective form factor determined from $e^+e^- \rightarrow p\bar{p}$, exhibiting a narrow peak or a very steep falloff around the $p\bar{p}$ mass threshold, which inspired many speculations and renewed the interests on the nucleon-antinucleon bound state.

X(1840) is a new structure discovered in the $J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$ process in 2013 with subdata sample of BESIII experiment, which is also located near the $p\bar{p}$ mass threshold. A further exploration of line shape of X(1840) is essential to have a better understanding of its nature. Therefore, the BESIII experiment performed an investigation on the $3(\pi^+ \pi^-)$ mass spectrum with 10 billion J/ψ events, which is about 45 times larger than the subdata sample used in the previous measurement.



The anomalous line shape of the resonant structure around $p\bar{p}$ mass threshold in $3(\pi^+\pi^-)$ mass spectrum. Credit: *Physical Review Letters* (2024). DOI: 10.1103/PhysRevLett.132.151901

An anomalous line shape of X(1840) near the $p\bar{p}$ mass threshold was observed for the first time. After many attempts, it was found that the model with a coherent sum of two Breit-Wigner parameterizations could provide a good description of data, which revealed a new resonance X(1880) with a [statistical significance](#) greater than 10σ , and the mass and width were determined to be $1882.1 \pm 1.7 \pm 0.7 \text{ MeV}/c^2$ and $30.7 \pm 5.5 \pm 2.4 \text{ MeV}/c$, respectively.

The proximity of its mass to the $p\bar{p}$ mass threshold supported the existence of a $p\bar{p}$ bound state. After publication, this result was selected as "Featured in Physics" by *Physical Review Letters*.

More information: M. Ablikim et al, Observation of the Anomalous Shape of X(1840) in $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ Indicating a Second Resonance Near $p\bar{p}$ Threshold, *Physical Review Letters* (2024). [DOI: 10.1103/PhysRevLett.132.151901](https://doi.org/10.1103/PhysRevLett.132.151901)

Provided by Chinese Academy of Sciences

Citation: Evidence of a new subatomic particle observed (2024, April 12) retrieved 2 May 2024 from <https://phys.org/news/2024-04-evidence-subatomic-particle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.