

## FASER measures high-energy neutrino interaction strength

April 8 2024, by Piotr Traczyk



Event displays identified by the FASER collaboration as candidates for an  $v_e$  (left) and a  $v_{\mu}$  (right) interacting in the detector. Invisible here, the neutrinos arrive from the left and then interact to create multiple tracks spraying out to the right (colored lines), one of which is identified as a charged lepton (labeled). Credit: FASER collaboration

Operating at CERN's Large Hadron Collider (LHC) since 2022, the FASER experiment is designed to search for extremely weakly interacting particles. Such particles are predicted by many theories beyond the Standard Model that are attempting to solve outstanding problems in physics such as the nature of dark matter and the matterantimatter imbalance in the universe.



Another goal of the experiment is to study interactions of high-energy neutrinos produced in the LHC collisions, particles that are nearly impossible to detect in the four big LHC experiments. Last week, at the annual Rencontres de Moriond conference, the FASER collaboration presented a measurement of the interaction strength, or "cross section," of electron neutrinos ( $v_e$ ) and muon neutrinos ( $v_\mu$ ).

This is the first time such a measurement has been made at a particle collider. Measurements of this kind can provide important insights across different aspects of physics, from understanding the production of "forward" particles in the LHC collisions and improving our understanding of the structure of the proton to interpreting measurements of high-energy neutrinos from astrophysical sources performed by neutrino-telescope experiments.

FASER is located in a side tunnel of the LHC accelerator, 480 meters away from the ATLAS detector collision point. At that location, the LHC beam is already nearly 10 meters away, bending away on its circular 27-kilometer path. This is a unique location for studying weakly interacting particles produced in the LHC collisions.

Charged particles produced in the collisions are deflected by the LHC magnets. Most neutral particles are stopped by the hundreds of meters of rock between FASER and ATLAS. Only very weakly interacting neutral particles like neutrinos are expected to continue straight on and reach the location where the detector is installed.

The probability of a neutrino interacting with matter is very small, but not zero. The type of interaction that FASER is sensitive to is where a neutrino interacts with a proton or a neutron inside the detector. In this interaction, the neutrino transforms into a charged "lepton" of the same family—an electron in the case of a  $v_e$ , and a muon in the case of a  $v_{\mu}$ —which is visible in the detector. If the energy of the neutrino is high,



several other particles are also produced in the <u>collision</u>.

The detector used to perform the measurement consists of 730 interleaved tungsten plates and photographic emulsion plates. The emulsion was exposed during the period from 26 July to 13 September 2022 and then chemically developed and analyzed in search of charged particle tracks.

Candidates for neutrino interactions were identified by looking for clusters of tracks that could be traced back to a single vertex. One of these tracks then had to be identified as a high-energy electron or muon.

In total, four candidates for an  $v_e$  interaction and eight candidates for a  $v_{\mu}$  interaction have been found. The four  $v_e$  candidates represent the first direct observation of electron neutrinos produced at a collider. The observations can be interpreted as measurements of neutrino interaction cross sections, yielding  $(1.2^{+0.9}_{-0.8}) \times 10^{-38} \text{ cm}^2 \text{ GeV}^{-1}$  in the case of the  $v_e$  and  $(0.5 \pm 0.2) \times 10^{-38} \text{ cm}^2 \text{ GeV}^{-1}$  in the case of the  $v_{\mu}$ .

The energies of the neutrinos were found to be in a range between 500 and 1700 GeV. No measurement of the neutrino interaction cross section had previously been made at energies above 300 GeV in the case of the  $v_e$  and between 400 GeV and 6 TeV in the case of the  $v_{\mu}$ .

The results obtained by FASER, posted to the *arXiv* preprint server, are consistent with expectations and demonstrate the ability of FASER to make neutrino cross-section measurements at the LHC. With the full LHC Run 3 data, 200 times more neutrino events will be detected, allowing much more precise measurements.

More information: First Measurement of the  $v_e$  and  $\nu_\mu$  Interaction



Cross Sections at the LHC with FASER's Emulsion Detector, *arXiv* (2024). DOI: 10.48550/arxiv.2403.12520

Provided by CERN

Citation: FASER measures high-energy neutrino interaction strength (2024, April 8) retrieved 21 May 2024 from <u>https://phys.org/news/2024-04-faser-high-energy-neutrino-interaction.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.