

Why figuring out how potassium is destroyed in stars is important to understanding the universe

April 16 2024, by Tracey Peake



Globular clusters like NGC 2419, visible in this image taken with the NASA/ESA Hubble Space Telescope, are not only beautiful, but also fascinating. Credit: ESA/Hubble & NASA

If you want to know where elements come from, look to the stars. Almost every element heavier than helium is formed through nuclear



reactions in stars. But which stellar processes are responsible for these elements? Can we find patterns in how much of each element we observe in different astrophysical environments, like stars, galaxies or globular clusters?

Recently, a team of NC State researchers focused on the process of potassium (K) destruction in globular clusters, looking at one cluster in particular: NGC 2419. The paper is <u>published</u> in the journal *Physical Review Letters*.

Globular clusters are groups of gravitationally bound stars. Astronomers have observed clear patterns in the relative amounts of different elements from star to star. One such pattern is between oxygen and sodium: stars within globular clusters that have more sodium have less oxygen, and vice-versa. This is known as the sodium-oxygen (Na-O) anticorrelation. Several other anticorrelations have also been discovered, which indicates that unique (sometimes unknown) processes occur in specific <u>globular clusters</u>.

In 2012, the first magnesium-potassium (Mg-K) anticorrelation was discovered in a specific globular cluster, called NGC 2419. An overall surplus of potassium was linked to reactions from hydrogen burning at temperatures between 80 and 260 million kelvin.

But the puzzling thing is that the stars in the cluster that showed the anticorrelation are relatively young, red giant stars. The cores of these stars should not be hot enough for <u>nuclear reactions</u> to alter the amount of Mg and K. The leading theory involved mixing with K and Mg from old stars in the cluster, but what has remained uncertain is the speed of the potassium-destroying reaction.

A research team attempted to recreate the potassium-destroying reaction by performing an experiment on a similar nuclear reaction (39 K + 3 He



 \rightarrow ⁴⁰Ca + d), at the Triangle Universities Nuclear Laboratory (TUNL).

This reaction is a proton-transfer reaction, where a proton from helium-3 (3 He) is transferred to potassium-39 (39 K), forming calcium-40 (40 Ca). This experimental reaction allows us to imitate the real reaction that occurs in a star where potassium is destroyed.

They found that not only can potassium be destroyed at <u>lower</u> <u>temperatures</u>, it is destroyed 13 times faster than previously thought at these temperatures.

The finding could change the way we model element creation in stars—not just for this specific case of NGC 2419, but also for other astrophysical models that include reactions on potassium.

More information: W. Fox et al, High Resolution Study of Ca40 to Constrain Potassium Nucleosynthesis in NGC 2419, *Physical Review Letters* (2024). DOI: 10.1103/PhysRevLett.132.062701. On arXiv: DOI: 10.48550/arxiv.2401.06754

Provided by North Carolina State University

Citation: Why figuring out how potassium is destroyed in stars is important to understanding the universe (2024, April 16) retrieved 17 May 2024 from <u>https://phys.org/news/2024-04-figuring-potassium-destroyed-stars-important.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.