

# A frozen chunk of genome rewrites our understanding of bird evolution

April 1 2024

---



A wompoo fruit-dove in Queensland, Australia. Unraveling a genetic mystery revealed that flamingos and doves are more distantly related than previously thought. Credit: Daniel J. Field

An enormous meteor spelled doom for most dinosaurs 65 million years ago. But not all. In the aftermath of the extinction event,

birds—technically dinosaurs themselves—flourished.

Scientists have spent centuries trying to organize and sort some 10,000 species of birds into one clear family tree to understand how the last surviving dinosaurs filled the skies. Cheap DNA sequencing should have made this simple, as it has for countless other species.

But birds were prepared to deceive us.

In a pair of new research papers released April 1 scientists reveal that another event 65 million years ago misled them about the true family history of birds. They discovered that a section of one chromosome spent millions of years frozen in time, and it refused to mix together with nearby DNA as it should have.



This section, just 2% of the bird genome, convinced scientists that most birds could be grouped into two major categories, with flamingos and doves as evolutionary cousins. The more accurate family tree, which accounts for the misleading section of the genome, identifies four main groups and identifies flamingos and doves as more distantly related.

"My lab has been chipping away at this problem of bird evolution for longer than I want to think about," said Edward Braun, Ph.D., the senior author of the [paper](#) published in the *Proceedings of the National Academy of Sciences* and a professor of biology at the University of Florida. "We had no idea there would be a big chunk of the genome that behaved unusually. We kind of stumbled onto it."

Braun supervised an international team of collaborators led by Siavash Mirarab, a professor of computer engineering at the University of California San Diego, to publish their evidence that this sticky chunk of DNA muddied the true history of bird evolution. Mirarab and Braun also contributed to a companion paper published in *Nature* that outlines the

updated bird family tree, which was led by Josefin Stiller at the University of Copenhagen.

Both papers are part of the [B10K avian genomics project](#) led by Guojie Zhang of Zhejiang University, Erich Jarvis of Rockefeller University, and Tom Gilbert of the University of Copenhagen.



Two mutually exclusive bird family trees. The top family tree lumps flamingos and doves, in blue and teal respectively, closely together, while the bottom

family tree does not. The top family tree was built around distortions in bird genomes that date back to the extinction of the dinosaurs. The bottom family tree is likely more accurate, after accounting for these genomic anomalies.

Credit: Edward Braun

[Ten years ago](#), Braun and his collaborators pieced together a family tree for the Neoaves, a group that includes the vast majority of bird species. Based on the genomes of 48 species, they split the Neoaves into two big categories: doves and flamingos in one group, all the rest in the other. When repeating a similar analysis this year using 363 species, a different [family tree](#) emerged that split up doves and flamingos into two distinct groups.

With two mutually exclusive family trees in hand, the scientists went hunting for explanations that could tell them which tree was correct.

"When we looked at the individual genes and what tree they supported, all of a sudden it popped out that all the genes that support the older tree, they're all in one spot. That's what started the whole thing," Braun said.

Investigating this spot, Braun's team noticed it was not as mixed together as it should have been over millions of years of sexual reproduction. Like humans, birds combine genes from a father and a mother into the next generation. But birds and humans alike first mix the genes they inherited from their parents when creating sperm and eggs. This process is called recombination, and it maximizes a species' genetic diversity by making sure no two siblings are quite the same.

Braun's team found evidence that one section of one bird chromosome had suppressed this recombination process for a few million years around the time the dinosaurs disappeared. Whether the extinction event

and the genomic anomalies are related is unclear.

The result was that the flamingos and doves looked similar to one another in this chunk of frozen DNA. But taking into account the full genome, it became clear that the two groups are more distantly related.

"What's surprising is that this period of suppressed recombination could mislead the analysis," Braun said. "And because it could mislead the analysis, it was actually detectable more than 60 million years in the future. That's the cool part."

Such a mystery could be lurking in the genomes of other organisms as well.

"We discovered this misleading region in birds because we put a lot of energy into sequencing [birds'](#) genomes," Braun said. "I think there are cases like this out there for other species that are just not known right now."

**More information:** Mirarab, Siavash, A region of suppressed recombination misleads neoavian phylogenomics, *Proceedings of the National Academy of Sciences* (2024). [DOI: 10.1073/pnas.2319506121](https://doi.org/10.1073/pnas.2319506121). [doi.org/10.1073/pnas.2319506121](https://doi.org/10.1073/pnas.2319506121)

Provided by University of Florida

Citation: A frozen chunk of genome rewrites our understanding of bird evolution (2024, April 1) retrieved 2 May 2024 from <https://phys.org/news/2024-04-frozen-chunk-genome-rewrites-bird.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.