


# Genetic underpinnings of environmental stress identified in model plant

April 11 2024



Transcriptome data under ABA-related stresses was analyzed using 216 pairs of *Arabidopsis* gene expression data accumulated in public databases. Genes with variable expression levels common to five stress treatments (ABA, salt, drought, osmotic pressure, and cold) were identified. Fourteen genes were upregulated in all stress treatments, and eight genes were downregulated in all treatments.

Credit: Mitsuo Shintani/Hiroshima University

Plants can be temperamental. Even weeds along the side of highways or pushing their way up in the cracks of concrete sidewalks can get stressed out by dehydration, cold, excess salt and more. Researchers at Hiroshima University have identified 14 genes that thale cress—a plant commonly used in genetic investigations since its genome is well documented—express more when responding to five specific stressors, as well as eight genes that the plant suppresses.

They [published](#) their results in the journal *Frontiers in Plant Science*.

"Abiotic stresses—as opposed to biotic stresses like pests or disease—such as drought, salinity and cold negatively affect plant growth and crop productivity. Understanding the [molecular mechanisms](#) underlying [plant responses](#) to these stressors is essential for [stress tolerance](#) in crops," said corresponding author Hidemasa Bono, professor in the Laboratory of Genome Informatics at Hiroshima University's Graduate School of Integrated Sciences for Life. Bono is also affiliated with the Laboratory of Bio-DX in the university's Genome Editing Innovation Center.

"The plant hormone abscisic acid (ABA) is significantly increased upon abiotic stressors, inducing physiological responses to adapt to stress and regulate [gene expression](#). Although many studies have examined the components of established stress signaling pathways, few have explored other unknown elements."

To better understand the molecular pathways that allow ABA to increase, the research team analyzed public RNA sequencing data on thale cress, or *Arabidopsis thaliana*. RNA sequencing is a technique that enables scientists to identify and quantify specific sequences of genetic instructions programmed in an organism's RNA. This data can reveal how different variables may increase or decrease the expression of certain genes.

Bono and his team specifically focused on five ABA-related stress conditions: ABA, when the hormone is applied directly to the plant; salt, which changes how the plant can use water; dehydration, or how much water the plant has; osmotic, when plant cells swell or shrink inappropriately; and cold.

"The data-driven studies have the advantage of analyzing large and independent datasets, which can lead to the identification of novel targets, distinct from the extensively studied established factors and accelerate the development of stress-tolerant crops," Bono said.

The researchers performed a [meta-analysis](#) of 216 paired datasets, combining those research results and reanalyzing them to identify where data might overlap or reveal previously unknown connections.

The meta-analysis revealed that 14 genes were commonly up-regulated and eight genes were commonly down-regulated across all five ABA-related stress responses investigated. Bono noted that some genes regulated by salt, dehydration and osmotic treatments were not regulated by ABA or cold stress, suggesting that they may be involved in the plant response through a different signaling pathway.

"Our meta-analysis revealed a list of candidate genes with unknown molecular mechanisms in ABA-dependent and ABA-independent stress responses," Bono said.

"These genes could be valuable resources for selecting genome editing targets and potentially contribute to the discovery of novel stress tolerance mechanisms and pathways in plants. We will continue to develop methods and utilize data from public databases and conduct [comparative analysis](#) from various angles to unravel the unknown mechanisms of stress response in plants."

**More information:** Mitsuo Shintani et al, Meta-analysis of public RNA sequencing data of abscisic acid-related abiotic stresses in *Arabidopsis thaliana*, *Frontiers in Plant Science* (2024). [DOI: 10.3389/fpls.2024.1343787](https://doi.org/10.3389/fpls.2024.1343787)

Provided by Hiroshima University

Citation: Genetic underpinnings of environmental stress identified in model plant (2024, April 11) retrieved 2 May 2024 from <https://phys.org/news/2024-04-genetic-underpinnings-environmental-stress.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.