

Ocean currents threaten to collapse Antarctic ice shelves, study finds

April 11 2024



Dotson Ice Shelf, Amundsen Sea, Antarctica. Credit: Taewook Park

A new study published in *Nature Communications* has revealed that the interplay between meandering ocean currents and the ocean floor induces upwelling velocity, transporting warm water to shallower depths. This mechanism contributes substantially to the melting of ice shelves in the Amundsen Sea of West Antarctica. These ice shelves are destabilizing rapidly and contributing to sea level rise.



Led by Taewook Park and Yoshihiro Nakayama, an international team of researchers from the Korea Polar Research Institute, Hokkaido University, and Seoul National University employed advanced ocean modeling techniques to investigate the underlying forces behind the rapid melting <u>ice shelves</u>.

In a departure from prior assumptions linking ice shelf melting primarily to winds over the Southern Ocean, this study underscores the significant role played by the interactions between meandering <u>ocean currents</u> and the ocean floor in driving the melting process.

The Pine Island and Thwaites ice shelves are among the fastest-changing in Antarctica and are of particular interest due to their vulnerability to warming ocean waters. They act as massive barriers restraining the glaciers behind them from flowing into the ocean.

However, their rapid melting and potential collapse pose a significant threat to coastal communities worldwide because of the resulting rise in global sea levels.

The study focused on the role of a layer of <u>warm water</u> beneath the frigid surface waters, known as the "modified Circumpolar Deep Water," in melting these ice shelves from below. "The intensity and trajectory of ocean currents encircling the ice shelves directly govern the influx of warm water, thereby intricately shaping their rate of melting," explains Taewook. This shows the importance of the ocean in understanding and addressing the impacts of climate change.

The researchers paid attention to the "thermocline depth," which is the depth of the interface between warmer deep waters and cooler surface waters. Variations in thermocline depth significantly affect the influx of warm water toward the ice shelves.



Until now, it has been believed that intensified westerly winds north of the Amundsen Sea propelled ocean currents along the shelf break, carrying warmer water toward ice shelf cavities. This phenomenon is particularly pronounced during El Niño events.

"Our findings challenge <u>conventional wisdom</u>," Nakayama says. "Our study underscores that the interplay between meandering ocean currents and the <u>ocean floor</u> generates upwelling velocity, bringing warm water to shallower depths. Subsequently, this warm water reaches the ice-ocean interface, accelerating ice shelf melting

"This internal oceanic process driving ice shelf melting introduces a novel concept. With this in mind, we have to reevaluate winds driving Antarctic ice loss, which can significantly impact future projections."

More information: Amundsen Sea Circulation Controls Bottom Upwelling and Antarctic Pine Island and Thwaites Ice Shelf Melting, *Nature Communications* (2024). DOI: 10.1038/s41467-024-47084-z

Provided by Hokkaido University

Citation: Ocean currents threaten to collapse Antarctic ice shelves, study finds (2024, April 11) retrieved 21 May 2024 from <u>https://phys.org/news/2024-04-ocean-currents-threaten-collapse-antarctic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.