


How phytochemical diversity affects plant adaptation to stress

April 16 2024, by Chen Na



Graphical abstract. Credit: *Science of The Total Environment* (2024). DOI: 10.1016/j.scitotenv.2024.172051

Phytochemical diversity plays a critical role in determining plant adaptation and fitness, as well as ecosystem functions and services. However, phytochemicals and their ecological adaptations have long been ignored. Fine (or absorptive) roots, microorganisms, and soil form the complex interface known as the rhizosphere. However, the variation in fine root chemistry along large-scale environmental gradients remains

poorly understood.

In a [study](#) published in *Science of the Total Environment*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) of the Chinese Academy of Sciences investigated the phytochemical [diversity](#) and its adaptation to abiotic and biotic stresses along different climatic gradients. By delving into the [chemical](#) constituents in fine roots, they sought to understand how these chemical constituents help plants cope with various environmental stresses.

The researchers aimed to explore the trends in the complexity of phytochemical composition (alpha diversity) and the variation in composition (beta diversity or endemism) and their relationships with microorganisms along environmental gradients. They measured the untargeted metabolomics of fine roots and the rhizosphere microbiome of 315 tree species along a macro-climatic [gradient](#) spanning tropical, subtropical, and subalpine region.

The results showed that the chemical diversity of subalpine tree species was higher than that of tropical tree species, but the variation in chemical diversity was smaller, which is conducive to adaptation to higher abiotic stress. Tropical tree species had higher variation in chemical diversity and chemical endemism, which is conducive to species coexistence and adaptation to complex biological stress.

In addition, there was extensive chemo-ecological niche differentiation among closely related [tree species](#) in all climate zones, and fine [root](#) chemical diversity was weakly influenced by phylogeny and mainly regulated by abiotic and biotic stresses.

"Our study is a positive exploration of phytochemical diversity patterns and their interactions with rhizosphere microorganisms along climatic gradients, which highlights the importance of future phytochemical

research," said Yang Jie of XTBG.

More information: Yazhou Zhang et al, Phytochemical diversity and their adaptations to abiotic and biotic pressures in fine roots across a climatic gradient, *Science of The Total Environment* (2024). [DOI: 10.1016/j.scitotenv.2024.172051](https://doi.org/10.1016/j.scitotenv.2024.172051)

Provided by Chinese Academy of Sciences

Citation: How phytochemical diversity affects plant adaptation to stress (2024, April 16)
retrieved 2 May 2024 from
<https://phys.org/news/2024-04-phytochemical-diversity-affects-stress.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.