

A powerful technique for tracking a protein's fleeting shape changes





IFS_{open} and IFS_{closed} conformational substates and identification of a kinetically trapped state. Credit: *Nature Structural & Molecular Biology* (2024). DOI: 10.1038/s41594-024-01260-3

Researchers at Weill Cornell Medicine have developed a powerful, new technique to generate "movies" of changing protein structures and speeds of up to 50 frames per second.



Senior author, Dr. Simon Scheuring, the Distinguished Professor of Anesthesiology Research at Weill Cornell Medicine and colleagues developed the new approach to gain a better understanding of how <u>biological molecules</u> change structurally over time.

Although investigators in this field routinely image static proteins and other molecules finely enough to resolve the positions of individual atoms, the resulting structural pictures or models are snapshots. Recording the dynamics of molecular structures—making movies—has been a much harder challenge. The lead author of the study is Yining Jiang, a doctoral candidate in the Weill Cornell Graduate School of Biomedical Sciences.

In their study, <u>published</u> April 17 in *Nature Structural & Molecular Biology*, the researchers used a relatively new measurement technique called high-speed atomic-force microscopy (HS-AFM), which employs an extremely sensitive probe to scan across molecules' surfaces, essentially feeling their structures. As a key innovation, the scientists found a method to isolate their target molecule, a single <u>protein</u>, thus avoiding effects from protein-to-protein interactions and enabling faster and more precise scanning.

The researchers applied their new single-molecule HS-AFM approach to a protein called Glt_{Ph} , a "transporter" that sits in the <u>cell membrane</u>, directing neurotransmitter molecules into the cell. Such transporters are among the favorite targets of structural biologists due to their complex and puzzling dynamics, and their importance in health and disease.

The researchers obtained dynamic structural data on Glt_{Ph} with an unprecedented combination of high spatial and time resolution—and stability, so that they could record tiny fluctuations in Glt_{Ph} 's structure continuously for minutes.





Workflow for the MEMPR method for HS-AFM study of membrane protein structure and dynamics. Credit: *Nature Structural & Molecular Biology* (2024). DOI: 10.1038/s41594-024-01260-3

An unsolved phenomenon in such proteins was termed "wanderlust" kinetics, meaning that molecules were reported to functionally change between high and low activity modes, for no obvious reason. The work revealed a previously unseen structural state of Glt_{Ph} , in which the transporter is locked and functionally asleep, uncovering the basis of 'wanderlust' kinetics.

The researchers emphasized that their new approach, which they are continually trying to optimize, is generalizable for studying other proteins, including membrane-embedded proteins. Overall, they said, this work opens up new possibilities to track the precise structure of a



protein moment-by-moment during its cycles of activity and rest.

More information: Yining Jiang et al, HS-AFM single-molecule structural biology uncovers basis of transporter wanderlust kinetics, *Nature Structural & Molecular Biology* (2024). DOI: 10.1038/s41594-024-01260-3

Provided by Weill Cornell Medical College

Citation: A powerful technique for tracking a protein's fleeting shape changes (2024, April 17) retrieved 21 May 2024 from <u>https://phys.org/news/2024-04-powerful-technique-tracking-protein-fleeting.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.