





A rechargeable nanotorch: Afterglow luminescence imaging tracks cell-based microrobots in real time

April 15 2024





- Long-life luminescence (> 10h)
- 660 nm-light charging

Rechargeable Nanotorch



Cell-based Microrobots

- Magnetically controlled
- Traceable



Credit: *Angewandte Chemie International Edition* (2024). DOI: 10.1002/anie.202400658

An afterglow luminescent nanoprobe opens up new possibilities for imaging living cells. As a research team [reports](#) in the journal *Angewandte Chemie International Edition*, their new "nanotorch" can continue to luminesce for more than 10 days after a single excitation.

This allows the routes taken through the body by microrobots to be tracked in real time. In addition, it can be "recharged" non-invasively with near-infrared (NIR) light in a non-contact manner.

Macrophages are important immune cells that "eat" bacteria as well as being involved in the disposal of cancer cells. In addition, they can take up drugs and transport them into cells, including tumor cells. If they take up [magnetic nanoparticles](#), macrophages can be guided by magnet to a target area within the body, such as a tumor. This allows macrophage "microrobots" to reduce the side effects associated with chemotherapy.

It would be useful to be able to track the microrobots over time as they move through the body. Fluorescence imaging techniques have been considered but require constant external irradiation. This causes is a high level of background noise resulting from the autofluorescence of many biomolecules. In addition, the limited penetration depth of the visible and UV light through tissues required into the tissue limits the depth of detection.

One alternative could be the use of probes that can be irradiated before the procedure and produce an afterglow. However, inorganic

nanoparticles with long-lasting afterglow harbor the risk that heavy metal ions will leak out; while [organic compounds](#) only luminesce for a short time and cannot be repeatedly excited.

A team from the Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences (China) collaborating with Koç University (Turkey) has now developed a "rechargeable nanotorch." It is made of multiple components: nanoparticles of a precursor to a luminescent organic molecule, photosensitizers (a hydrophobic analog of methylene blue), and [polyethylene glycol](#) equipped with cell-penetrating peptides.

The photosensitizer absorbs NIR light and excites surrounding oxygen molecules. This highly reactive singlet oxygen then binds to the precursor and forms a dioxetane group, a four-membered ring made of two oxygen and two carbon atoms. This undergoes a rearrangement that releases the desired luminescent molecule and emits excess energy by luminescing. After the initial irradiation, the nanotorches continue to luminesce for ten days.

Once depleted, the nanotorches can be "remotely" recharged and made to luminesce again by external radiation with NIR light, which can penetrate deep into tissues—multiple times. This requires the relative amounts of photosensitizer and luminescent molecule precursor to be selected so that only some of the precursors are activated with each irradiation. This allows for imaging over longer periods of time.

The China team led by Pengfei Zhang, Ping Gong and Lintao Cai collaborated with the Turkey team led by Safacan-Kolemen to introduce these new nanotorches into macrophage-based microrobots and were able to follow their magnet-guided path through the bodies of mice in real time through the luminescence signals.

More information: Gongcheng Ma et al, Rechargeable Afterglow

Nanotorches for In Vivo Tracing of Cell-Based Microrobots,
Angewandte Chemie International Edition (2024). DOI:
[10.1002/anie.202400658](https://doi.org/10.1002/anie.202400658)

Provided by Wiley

Citation: A rechargeable nanotorch: Afterglow luminescence imaging tracks cell-based microrobots in real time (2024, April 15) retrieved 2 May 2024 from <https://phys.org/news/2024-04-rechargeable-nanotorch-afterglow-luminescence-imaging.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.