

Why we need to rethink what we know about dust

April 2 2024, by Adrian Chappell



Credit: CC0 Public Domain

You may think of dust as an annoyance to be vacuumed and disposed of, but actually, on a grander scale, it is far more important than most people realize. Globally, dust plays a critical role in regulating our

climate, radiation balance, nutrient cycles, soil formation, air quality and even human health.

But our understanding of it has been hampered by limitations in current mathematical models. These models, built on methods developed decades ago, [struggle](#) to accurately simulate the properties and quantities of dust.

The latest [research](#) by my colleagues and I sheds light on these limitations and suggests a more nuanced picture of dust. Our [findings](#) reveal that dust emissions are not constant but shift seasonally and between hemispheres, across deserts and shrublands. This challenges the long-held [notion](#) that north Africa and the Middle East are the dominant sources of global dust.

Using two types of satellite data, our research suggests that dust emissions during [dust storms](#) are rare and localized, much like lightning strikes, and occur in constantly shifting locations.

Dust's complexity

The [cycle of dust emission](#), transport and deposition has positive and negative effects on our environment. Nutrients in deposited dust fertilize our oceans and rainforests. But dust from eroded sediment can also damage plants and trees and disrupt photosynthesis, while dust deposited on ice increases the speed at which it melts.

Variations in dust composition, like mineral type and color, create a complex cocktail of particles injected into the atmosphere. This, in turn, interacts with clouds to influence how sunlight is reflected or absorbed, ultimately regulating Earth's temperature.

So, it is vital we have an accurate understanding of where dust emissions

are coming from, in what quantities, how dust is transported across the planet and where it ends up.

Dust emission models were developed nearly 30 years ago when there was far less data available. Consequently, those now classical dust cycle models made some assumptions. One important assumption was that Earth's land surface was uniformly covered in perpetually loose and dry material, which was always available and caused [dust emissions](#).

However, we now know from [field measurements](#) that soils are often crusted or covered in different types of gravel. The threshold for the wind to lift the soil and release it to the atmosphere was also assumed to be fixed and unchanging over time.

We also now know that sediment moves around the landscape and may not always be available. Vegetation covering the soil reduces the wind's speed at reaching the soil surface, which then reduces dust emission. Dust models still assume that "greenness" indicates the presence of vegetation. However, in drylands where most dust emission happens, the vegetation is often brown, but its roughness still reduces the wind's speed and shelters the soil from dust emission.

Consequently, classical dust cycle models have over-estimated the amount of dust emission. These weaknesses have remained since models were developed. This is mainly because modelers assume that by adjusting their dust cycle models to the measurements of dust in the atmosphere they overcome any weaknesses in the dust emission modeling.

A new approach

Almost a decade ago we developed a new approach using [shadow](#) to estimate how much of the wind's speed is reduced by roughness, such as

vegetation, on the Earth's surface. This approach was still limited by the previous model assumptions described.

However, during the pandemic, traditional field studies became impossible. So, we adopted a new approach. Using satellites, we produced a global collection of dust emission points. This provided valuable data and paved the way for further [research](#).

We found that existing models overestimated the role of north Africa as the primary source of global dust emissions. Our research shows that [dust emissions](#) shift seasonally and between hemispheres, from deserts in east Asia, the Middle East and north Africa as well as shrublands in Australia and North America.

Current models have only been providing a fraction of the story based on dust in the atmosphere above north Africa and the Middle East. Little dust emission was predicted to occur in the southern hemisphere. But this contrasts with field observations and the experiences of people in those regions.

These new findings are crucial for large scale models because the properties of dust are different depending on where they come from. Not only that, but dust may change as it is transported within a hemisphere to different destinations where it settles on land, in our oceans and on icecaps.

Our new understanding of dust distribution, quantity and seasonal shifts has significant implications. It will require revisions to historical reconstructions that explain past climate changes. Our findings will also influence future climate projections and how the dust cycle interacts with the carbon, energy and water cycles of Earth's systems.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Why we need to rethink what we know about dust (2024, April 2) retrieved 21 May 2024 from <https://phys.org/news/2024-04-rethink.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--