


Research reveals a surprising topological reversal in quantum systems

April 18 2024, by Oliver Morsch



The results of the ETH researchers as an homage to Andy Warhol. The image shows the experimental results of topological pumping. Credit: Quantum Optics Group / ETH Zurich

In principle, one shouldn't compare apples to oranges. However, in topology, which is a branch of mathematics, one must do just that. Apples and oranges, it turns out, are said to be topologically the same since they both lack a hole—in contrast to doughnuts or coffee cups, for instance, which both have one (the handle in the case of the cup), and

thus are topologically equal.

In a more abstract way, [quantum systems](#) in physics can also have a specific apple or doughnut topology, which manifests itself in the energy states and motion of particles. Researchers are very interested in such systems as their topology makes them robust against disorder and other disturbing influences, which are always present in natural physical systems.

Things get particularly interesting if in addition, the particles in such a system interact, meaning that they attract or repel each other, like electrons in solids. Studying topology and interactions together in solids, however, is extremely difficult. A team of researchers at ETH led by Tilman Esslinger has now managed to detect topological effects in an artificial solid, in which the interactions can be switched on or off using magnetic fields.

Their results, which [appear](#) in *Science*, could be used in quantum technologies in the future.

Transport by topology


Zijie Zhu, a Ph.D. student in Esslinger's lab and first author of the study, and his colleagues constructed the artificial solid using extremely [cold atoms](#) (fermionic potassium atoms), which were trapped in spatially periodic lattices using laser beams. Additional [laser beams](#) caused the energy levels of adjacent lattice sites to move up and down periodically, out of sync with respect to each other.

After some time, the researchers measured the positions of the atoms in the lattice, initially without interactions between the atoms. In this experiment they observed that the doughnut topology of the energy states caused the particles to be transported by one lattice site, always in

the same direction, at each repetition of the cycle.

"This can be imagined as the action of a screw," says Konrad Viebahn, Senior Postdoc in Esslinger's team. The screwing motion is a clockwise rotation around its axis, but the screw itself moves in the forward direction as a result. With each revolution the screw advances a certain distance, which is independent of the speed at which one turns the screw. Such a behavior, also known as topological pumping, is typical of certain topological systems.

But what if the screw hits an obstacle? In the experiment of the ETH researchers, that obstacle was an additional laser beam that restricted the freedom of movement of the atoms in the longitudinal direction. After around 100 revolutions of the screw, the atoms ran into a wall, as it were. In the analogy used above, the wall represents an apple topology in which topological pumping cannot take place.



Using laser beams (yellow and red) the researchers create a lattice, in which the atoms are trapped (left) and transported through the lattice by topological pumping (right). Credit: Quantum Optics Group / ETH Zurich

Surprising return

Surprisingly, the atoms didn't simply stop at the wall, but suddenly turned around. The screw was thus moving backwards, although it kept being turned clockwise. Esslinger and his team explain this return by the two doughnut topologies that exist in the lattice—one with a clockwise-turning doughnut and another one that turns in the opposite direction. At the wall, the atoms can change from one [topology](#) to the other, thus inverting their direction of motion.

Then the researchers switched on a repulsive interaction between the atoms and watched what happened. Again, they were in for a surprise: The atoms now turned around at an invisible barrier even before reaching the laser wall.

"Using model calculations, we were able to show that the invisible barrier was created by the atoms themselves through their mutual repulsion," explains Ph.D. student Anne-Sophie Walter.

Qubit highway for quantum computers

"With these observations we have taken a big step towards a better understanding of interacting topological systems," says Esslinger, who studies such effects. As a next step, he wants to perform further experiments to investigate whether the topological screw is as robust as expected with respect to disorder, and how the atoms behave in two or three spatial dimensions.

Esslinger also has some practical applications in mind. For instance, the transport of atoms or ions by topological pumping could be used as a qubit highway to take the qubits (quantum bits) in quantum computers to the right places without heating them up or disturbing their quantum states.

More information: Zijie Zhu et al, Reversal of quantized Hall drifts at noninteracting and interacting topological boundaries, *Science* (2024).

[DOI: 10.1126/science.adg3848](https://doi.org/10.1126/science.adg3848)

Provided by ETH Zurich

Citation: Research reveals a surprising topological reversal in quantum systems (2024, April 18) retrieved 22 May 2024 from <https://phys.org/news/2024-04-reveals-topological-reversal-quantum.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--