


Unlocking the secrets of disease resistance in chrysanthemums: A holistic approach to combating black spot disease

April 3 2024



Differences in disease phenotype of different plants after inoculation. Credit: *Ornamental Plant Research*

Chrysanthemum, celebrated for its ornamental, medicinal, and beverage value, faces significant threats from bacterial and fungal infections, particularly black spot disease caused by *Alternaria alternata*, which leads to severe economic losses. Current research has focused on developing resistant germplasm as an eco-friendly alternative to pesticides.

Despite these efforts, the search for germplasm with exceptional [disease resistance](#) is ongoing, and the integration of physical and chemical plant defenses and [secondary metabolites](#) into breeding strategies remains underexplored. Therefore, further investigation into how these defenses contribute to resistance against black spot disease in chrysanthemum is of great importance to enhance breeding efforts for disease-resistant cultivars.

Ornamental Plant Research has recently [published](#) a paper titled "Genetic resources resistant to black spot (*Alternaria alternata*) identified from Chrysanthemum-related genera and potential underlying mechanisms."

In this study, researchers employed simplified detached leaf inoculation assay and whole plant inoculation assay to assess the resistance of 14 chrysanthemum-related genera (CRG) to black spot disease. After artificial inoculation and identification based on disease indices, two disease-resistant [germplasm](#) resources (R), 11 moderately resistant materials (MR), and one sensitive (S) material were obtained.

The results showed good reproducibility and correlation between methods, highlighting *C. japonense* and *A. parviflora* as resistant, and revealing significant physical and chemical defense mechanisms contributing to disease resistance.

Further analysis focused on the leaf epidermis structure of resistant and susceptible germplasms, revealing notable differences in trichome

morphology and density, stomatal characteristics, and wax content, which correlated with resistance levels. For instance, R1 had significantly higher trichome density and wax content, contributing to its resistance. Additionally, [volatile organic compounds](#) (VOCs) analysis indicated that resistant germplasms produced higher terpenoid content, with Germacrene D significantly inhibiting *A. alternata* growth, suggesting its role in chemical defense.

In conclusion, the study not only identified *C. japonese* and *A. parviflora* as resistant through a combination of physical (trichome density, stomatal closure, wax content) and chemical defenses (high VOCs and terpenoid content) but also demonstrated the importance of both types of defenses in disease resistance. These findings enhance our understanding of CRG resistance to black spot disease, offering insights into breeding strategies and the potential use of physical and chemical traits as markers for disease resistance.

More information: Qingling Zhan et al, Genetic resources resistant to black spot (*Alternaria alternata*) identified from Chrysanthemum-related genera and potential underlying mechanisms, *Ornamental Plant Research* (2023). [DOI: 10.48130/opr-0023-0023](https://doi.org/10.48130/opr-0023-0023)

Provided by Maximum Academic Press

Citation: Unlocking the secrets of disease resistance in chrysanthemums: A holistic approach to combating black spot disease (2024, April 3) retrieved 16 May 2024 from <https://phys.org/news/2024-04-secrets-disease-resistance-chrysanthemums-holistic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.