


# Study sheds new light on cross-species virus spillovers that can cause pandemics

April 23 2024



Guano consumption by wildlife in Budongo Forest Reserve, Uganda. Credit: *Communications Biology* (2024). DOI: 10.1038/s42003-024-06139-z

A study led by the University of Stirling jointly with the University of Wisconsin-Madison has shed new light on cross-species virus spillovers

that can cause pandemics.

A researcher from the Faculty of Natural Sciences at Stirling discovered animals in an African forest eating bat excrement known as [guano](#) after a key food source in the region disappeared following selective deforestation. The paper "Selective deforestation and exposure of African wildlife to bat-borne viruses" was [published](#) in the journal *Communications Biology*.

Lab analysis of the bat guano identified a range of viruses, including a betacoronavirus related to SARS-CoV-2, the [virus](#) that caused the COVID-19 pandemic. It remains unknown whether the betacoronavirus found in the guano is transmissible to humans, but it does offer an example of how new infections might jump species barriers.

The study was prompted when Dr. Pawel Fedurek observed wild chimpanzees consume bat guano from a tree hollow in Budongo Forest, Uganda.

Dr. Fedurek then set up cameras which recorded chimps, monkeys and antelope eating the excrement.

The guano is an alternative source of crucial minerals after the palm trees the mammals once consumed were harvested to extinction locally. The palm was used by people in Budongo to dry tobacco leaves which are then sold to international companies.

Scientists do not yet fully understand the earliest stages of virus spillovers, which can lead to the deaths of tens of thousands of people, but they are thought to involve complex causal chains that begin with humans altering the environment.

Dr. Pawel Fedurek, an expert in animal behavior at the University of

Stirling, said, "Our research illustrates how a subtle form of selective deforestation, ultimately driven by a global demand for tobacco, can expose wildlife and, by extension, humans to viruses residing in bat guano, increasing virus spillover risk.

"Studies like ours shed light on the triggers and pathways of both wildlife-to-wildlife and wildlife-to-human virus transmission, ultimately improving our abilities to prevent outbreaks and pandemics in the future.

"Most research into outbreaks and pandemics has focused on curtailing the spread of the virus, by finding an effective vaccine for example, rather than preventing animal-to-human virus transmission from happening in the first place. Our work stresses the importance of studying disease ecology before diseases enter humans."

After discovering mammals consuming guano, Dr. Fedurek and the project co-lead Dr. Caroline Asiimwe (then Conservation Coordinator of the Budongo Conservation Field Station, Uganda) involved Professor Tony Goldberg (University of Wisconsin-Madison, U.S.), who is a world-renowned expert in epidemiology and evolution of infectious diseases.

Professor Goldberg identified the viruses in the guano in collaboration with researchers who quantified the mineral content of the guano (Professor Jessica Rothman, Hunter College of the City University of New York, U.S.) and established the probability of the novel betacoronavirus to infect the three [mammalian species](#) and humans (Gregory K. Rice, Naval Medical Research Command, U.S.).

Due to the magnitude and complexity of the laboratory analyses involved, the entire project took around six years to complete.

## **Textbook example**

Professor Goldberg said, "Our study links tobacco farming to exposure of at least three species of African wildlife to viruses of bats. We believe this is a textbook example of how new infections might jump species barriers even before they get into humans.

"This is important since the initial stages of the onset of outbreaks, epidemics and pandemics have been particularly elusive for science.

"We may have stumbled upon a series of events that is usually hidden from the view of epidemiologists and public health officials. Our study might be particularly relevant to the origins of bat-borne disease such as coronaviruses and, perhaps, Ebola."

It is hoped the discovery may make it possible to enact interventions that break these sorts of causal chains, ultimately helping prevent future pandemics.

## Health consequences

Dr. Asiimwe said, "Our study demonstrates that human activities that alter the environment can lead to devastating health consequences to both wildlife and humans and thus we should urgently learn to use natural resources sustainably.

"Considerable research efforts are also needed to investigate how forest degradation, and other forms of human activity, can affect wildlife's behavior in a way that exposes them and humans to dangerous viruses."

**More information:** Pawel Fedurek et al, Selective deforestation and exposure of African wildlife to bat-borne viruses, *Communications Biology* (2024). [DOI: 10.1038/s42003-024-06139-z](https://doi.org/10.1038/s42003-024-06139-z)

Provided by University of Stirling

Citation: Study sheds new light on cross-species virus spillovers that can cause pandemics (2024, April 23) retrieved 22 May 2024 from <https://phys.org/news/2024-04-species-virus-spillovers-pandemics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.