


New technique lets scientists create resistance-free electron channels

April 9 2024, by Theresa Duque



Scanning tunneling microscopy image of a chiral interface state wavefunction (bright stripe) in a quantum anomalous Hall insulator made from twisted monolayer-bilayer graphene. Credit: Canxun Zhang/Berkeley Lab

An international research team led by Lawrence Berkeley National Laboratory (Berkeley Lab) has taken the first atomic-resolution images and demonstrated electrical control of a chiral interface state—an exotic

quantum phenomenon that could help researchers advance quantum computing and energy-efficient electronics.

The chiral interface state is a conducting channel that allows electrons to travel in only one direction, preventing them from being scattered backward and causing energy-wasting [electrical resistance](#). Researchers are working to understand better the properties of chiral interface states in real materials, but visualizing their spatial characteristics has proved to be exceptionally difficult.

But now, for the first time, atomic-resolution images captured by a research team at Berkeley Lab and UC Berkeley have directly visualized a chiral interface state. The researchers also demonstrated the on-demand creation of these resistance-free conducting channels in a 2D insulator.

Their work, which was reported in the journal [Nature Physics](#), is part of Berkeley Lab's broader push to advance [quantum computing](#) and other quantum information system applications, including the design and synthesis of quantum materials to address pressing technological needs.

"Previous experiments have demonstrated that chiral interface states exist, but no one has ever visualized them with such high resolution. Our work shows for the first time what these 1D states look like at the atomic scale, including how we can alter them—and even create them," said first author Canxun Zhang, a former graduate student researcher in Berkeley Lab's Materials Sciences Division and the Department of Physics at UC Berkeley. He is now a postdoctoral researcher at UC Santa Barbara.

Chiral interface states can occur in certain types of 2D materials known as quantum anomalous Hall (QAH) insulators that are insulators in bulk but conduct electrons without resistance at one-dimensional

"edges"—the physical boundaries of the material and interfaces with other materials.

To prepare chiral interface states, the team worked at Berkeley Lab's Molecular Foundry to fabricate a device called twisted monolayer-bilayer graphene, which is a stack of two atomically thin layers of graphene rotated precisely relative to one another, creating a moiré superlattice that exhibits the QAH effect.

In subsequent experiments at the UC Berkeley Department of Physics, the researchers used a [scanning tunneling microscope](#) (STM) to detect different electronic states in the sample, allowing them to visualize the wavefunction of the chiral interface state. Other experiments showed that the chiral interface state can be moved across the sample by modulating the voltage on a gate electrode placed underneath the graphene layers.

In a final demonstration of control, the researchers showed that a voltage pulse from the tip of an STM probe could "write" a chiral interface state into the sample, erase it, and even rewrite a new one where electrons flow in the opposite direction.

The findings may help researchers build tunable networks of electron channels with promise for energy-efficient microelectronics and [low-power](#) magnetic memory devices in the future, and for quantum computation making use of the exotic electron behaviors in QAH insulators.

The researchers intend to use their technique to study more exotic physics in related materials, such as anyons, a new type of quasiparticle that could enable a route to quantum computation.

"Our results provide information that wasn't possible before. There is

still a long way to go, but this is a good first step," Zhang said.

More information: Canxun Zhang et al, Manipulation of chiral interface states in a moiré quantum anomalous Hall insulator, *Nature Physics* (2024). [DOI: 10.1038/s41567-024-02444-w](https://doi.org/10.1038/s41567-024-02444-w)

Provided by Lawrence Berkeley National Laboratory

Citation: New technique lets scientists create resistance-free electron channels (2024, April 9) retrieved 17 May 2024 from <https://phys.org/news/2024-04-technique-scientists-resistance-free-electron.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--